

prevent undesirable taste and discoloration. In the USSR, the recommended maximum permissible concentration of  $\text{KMnO}_4$  is 0.1 mg/l (as Mn). The recommendation is intended to prevent the discoloration of water by manganese (Shigan and Vitvitskaya, 1971).

For marine waters, the U.S. EPA (1976) has recommended a criterion for manganese of 0.1 mg/l for the protection of consumers of marine mollusks. Although the rationale for this criterion is not detailed, it is partially based on the observation that manganese can bioaccumulate by "factors as high as 12,000" in marine mollusks.

## 9.2. SUMMARY OF TOXICITY

Manganese is an essential element for humans and animals. The concentration of manganese present in individual tissues, particularly in the blood, is controlled after ingestion by homeostatic mechanisms and remains remarkably constant in spite of rapid fluctuations in intake (Cotzias, 1958). The main routes of absorption are the gastrointestinal and respiratory tracts. Acute poisoning by manganese may occur in exceptional circumstances where large amounts of manganese compounds are ingested or inhaled. Freshly formed manganese oxide fumes of respirable particle size can cause metal fume fever but are not believed to cause permanent damage (Piscator, 1976). The most pronounced toxic effects of manganese are a CNS syndrome known as chronic manganese poisoning (manganism) and manganese pneumonitis.

The adverse effect on the CNS begins with a psychiatric disturbance followed by a neurologic phase resembling Parkinson's disease. Manganism has been well described in the literature with clinical details for case clusters (Flinn et al., 1940; Penzler, 1955; Rodier, 1955; Chandra et al., 1974). Cotzias (1962) described three phases -- a prodromal phase with insidious onset including psychiatric disturbances, the extrapyramidal

disease with symptoms of awkward speech and loss of skilled movement, and typical manganism with severe rigidity, tremor, and inappropriate emotional reactions.

Manganism has been reported in workers in ore crushing and packing mills, in the production of ferroalloys, in the use of manganese alloys in the steel industry and in the manufacture of dry cell batteries and welding rods. Very high concentrations of manganese have been found in mines where cases of manganism were reported. The manganese air concentration in the immediate vicinity of rock drilling in Moroccan mines was  $\sim 450 \text{ mg/m}^3$  in one mine and  $\sim 250 \text{ mg/m}^3$  in another (Rodier, 1955). In two reports from Chilean mines (Ansola et al., 1944a,b; Schuler et al., 1957) the air concentrations of manganese varied from  $62.5\text{-}250 \text{ mg/m}^3$  and from  $0.5\text{-}46 \text{ mg/m}^3$ , respectively.

In ferromanganese factories, neurological and psychic disturbances indicating manganese poisoning have been observed at manganese levels as low as  $2\text{-}5 \text{ mg/m}^3$  of air (Suzuki et al., 1973a,b).

While manganism and its association with manganese has been well described, a dose-response relationship in man cannot be evaluated because duration of exposure is not well documented. Also, early signs of the disease were sought in only a few studies (Saric et al., 1977; Tanaka and Lieben, 1969), and none of the reported studies employed a standard cohort design (e.g., there was no follow-up or comprehensive characterization of the exposed populations).

A high incidence of pneumonia and other respiratory ailments has been reported in workers with occupational exposure to manganese (Baader, 1937; Lloyd-Oavies, 1946; Rodier, 1955; Saric, 1978) and in inhabitants living around factories manufacturing ferromanganese or manganese alloys (Elstad,

1939; Suzuki, 1970). The increased incidence of pulmonary disease found in exposure to low concentrations of manganese is not necessarily directly attributable to manganese itself. Manganese exposure may increase susceptibility to pneumonia or other acute respiratory diseases by disturbing the normal mechanism of lung clearance. Some investigators have suggested that long-term exposure to manganese may contribute to the development of chronic lung disease (Saric and Lucic-Palaic, 1977), but there is little data to demonstrate this conclusively, particularly at ambient levels.

Effects on the cardiovascular system include reports of decreased systolic blood pressure in humans occupationally exposed via inhalation (Saric and Hrustic, 1975). This symptom was also shown to occur experimentally in orally exposed rats (Kimura et al., 19... Reports about the effects of manganese on human blood and hemoglobin show conflicting results that have not been resolved by animal studies. The studies are difficult to compare because of variations in exposure and in the severity of the effect.

There have been reports of impotence in a majority of workers affected by manganese (Chandra et al., 1974; Emara et al., 1971; Rodier, 1955; Penalver, 1955). There is some experimental evidence of reproductive effects in laboratory animals. Degenerative changes in the testes of rats have been produced by excessive levels of manganese administered by multiple intraperitoneal injections (Chandra, 1971; Singh et al., 1974, 1975; Chandra et al., 1975; Tandon et al., 1975; Shukla and Chandra, 1977) and by single intratracheal injections in rabbits (Chandra et al., 1973a). Chronic dietary exposure to manganese has caused decreased organ weight for the preputial gland, seminal vesicle and testis in mice (Gray and Laskey, 1980), and decreased serum testosterone levels and reduced pregnancy percentage in rats (Laskey et al., 1982).

Manganese dichloride increased the incidence of lymphosarcomas in DBA/1 mice following twice weekly subcutaneous or intraperitoneal injections for 6 months (DiPaolo, 1964), and elicited slightly elevated tumor incidence in a Strain A mouse lung tumor bioassay (Stoner et al., 1976). Single or repeated intramuscular injections of  $MnO_2$  or manganese powder did not result in increased incidences of lymphosarcomas, leukemias or local sarcomas in either sex of F344 rats or female Swiss mice. However, repeated intramuscular injections of an organomanganese compound, MAA, elicited statistically significant increases in injection site tumors in both sexes of F344 rats (Furst, 1978). Oral administration of manganese powder for 12 months twice monthly did not induce lymphomas and/or leukemias or fibromas in either sex of F344 rats (Furst, 1978). Although the results of the studies with divalent manganese are suggestive of carcinogenic activity, non-natural routes of administration were employed.

Some reported animal studies imply a carcinogenic potential for manganese, but the data are inadequate to support this conclusion (i.e., local injection site sarcomas in F344 rats, a marginal response in Strain A mice, and inadequate data in the experiment with DBA/1 mice). No epidemiologic information relating manganese exposure to cancer occurrence in humans has been located. Using IARC criteria (IARC, 1980), the available evidence for manganese carcinogenicity would be rated inadequate for animals and "no data available" for humans (Group 3). Consequently, the documented toxic effects are of more practical concern.

### 9.3. SPECIAL GROUPS AT RISK

Several researchers have mentioned the marked differences in individual susceptibility to inhaled manganese (Rodier, 1955; Penalver, 1955; Cotzias,

1958). They speculated that this may have been caused by alcoholism, syphilis, carbon monoxide, lesions of the excretory system, or the physiological or pathological condition of the respiratory tract. While it is reasonable to assume that an individual with an impaired ability to clear inhaled manganese or to excrete absorbed manganese would be at increased risk of adverse effects, no studies exist to confirm this.

Experimental studies suggest that populations at greatest risk of adverse effects due to manganese exposure are the very young and those with iron deficiency. The evidence for increased absorption and retention of manganese occurring in iron deficiency was shown in an inhalation study in humans (Mena et al., 1969, 1974), dietary studies in humans (Thomson et al., 1971), and ingestion studies in experimental animals (Rehnberg et al., 1982; Kostial et al., 1980).

Ingestion studies give useful information on the effects of inhalation exposures because most inhaled manganese is cleared to the gastrointestinal tract. The early neonatal period may be critical for metal accumulation because the very young also have an increased intestinal absorption and retention of manganese. This has been demonstrated in preweanling mice and rats (Kostial et al., 1978; Rehnberg et al., 1980) and in infants (Mena et al., 1974). High retention of manganese in the tissues, particularly the liver and brain, is associated with the limited excretion of manganese in the preweanling rat (Miller et al., 1975).

Kostial et al. (1978) report that oral toxicity measured by LD<sub>50</sub> values is greater in very young (2 weeks) as well as old (54 weeks) rats, but not as high as expected based on the rate of intestinal absorption. Although the neonate has not been shown to have increased sensitivity to metals, the early accumulation of manganese must be considered as an additional risk factor.

Another population at high risk is workers exposed to manganese at or near the recommended TLV. Because neurological symptoms have been reported at concentrations below this limit, the TLV of  $5 \text{ mg/m}^3$  has a low margin of safety.

#### 9.4. EFFECTS OF MAJOR CONCERN AND EXPOSURE/RESPONSE INFORMATION

9.4.1. Effects of Major Concern. The key health effects of manganese are in the CNS and the lungs. The effect on the CNS, manganism, is irreversible and severely incapacitating although not directly associated with lethality. The pulmonary effects reported at levels below  $1 \text{ mg/m}^3$  are for the most part reversible but can limit function or impose disability such as increased wheezing, bronchitis, or increased susceptibility to respiratory illness. The lowest reported exposure levels associated with life threatening diseases such as pneumonia have been similar to ranges associated with chronic manganese poisoning, 0.3 and up for brain effects.

Several endpoints suggested as effects from exposure to manganese are nonspecific, inconclusive or lack documentation in humans, such as degenerative changes in the testes, or decreased blood pressure. Sexual dysfunction has often been reported as an early effect of manganism at levels associated with other effects on the CNS.

9.4.2. Exposure/Response Information. Tables 6-1 and 6-3 show that human exposure to levels below  $5 \text{ mg/m}^3$  has been associated with adverse effects to the CNS. These effects are either advanced manganism or a constellation of signs indicative of early stages of the disease (Suzuki et al., 1973a; Chandra et al., 1981). There is some evidence suggesting that exposure to levels below  $1 \text{ mg/m}^3$  is associated with nonspecific symptoms which are common in early manganism and with abnormal neurological findings. However, in studies at these levels the findings reported could not be definitively attributed to manganese exposure (Saric et al., 1977; Chandra et al., 1981).

Studies of respiratory effects in humans (summarized in Table 6-8) show pulmonary system adverse effects at levels below  $1 \text{ mg/m}^3$ . Schoolchildren exposed to manganese emissions estimated at  $\sim 3\text{--}11 \text{ } \mu\text{g/m}^3$  from a ferro-manganese plant developed an increase in respiratory symptoms compared with controls such as sore throat, wheezing and sputum on arising (Nogawa et al., 1973). Saric and Lucic-Palaic (1977) reported increased chronic bronchitis in workers exposed to  $0.4\text{--}16 \text{ mg/m}^3$  but the results at ambient levels  $<1 \text{ mg/m}^3$  (Saric et al., 1975) were inconclusive because no exposure-response relationship was seen and confounding factors were not controlled.

There are many pulmonary endpoints that vary according to level of exposure. Although exposure ranges are so broad that the exposure/response relationship is sometimes masked, a continuum of effects has been observed which qualitatively supports the pulmonary endpoint. Pulmonary effects reported and supported include pneumonia (Elstad, 1939; Lloyd-Davies, 1946; Wassermann and Mihail, 1961), chronic bronchitis (Lloyd-Davies, 1946; Saric and Lucic-Palaic, 1977), radiographic changes and fibrosis (Wassermann and Mihail, 1961) and airways disability (Nogawa et al., 1973).

Animal studies also qualitatively support the association between pulmonary effects and manganese exposure. Table 9-1 summarizes the animal studies of the adverse effects of manganese inhalation. Pathological changes and decreased resistance to infection occur in a variety of species at levels above  $0.7 \text{ mg/m}^3$ . Inflammatory reactions and decreased resistance to infection have been observed in mice (Malgetter et al., 1976; Adkins et al., 1980c). Nishiyama et al. (1975) report pulmonary congestion and inflammatory changes in mice and monkeys after 5 months exposure to  $3 \text{ mg/m}^3$  and less severe changes at  $0.7 \text{ mg/m}^3$ . Suzuki et al. (1978) describe radiologic changes after 10 months of exposure to 3 and  $0.7 \text{ mg/m}^3$ .

TABLE 9-1

Studies of Manganese Inhalation in Animals -- Summary of Effect Levels

Species	Duration	Concentration (mg/m <sup>3</sup> )	System Examined	Effects*	Reference
MnO <sub>2</sub>					
Mouse	14 days	0.7	pulmonary	pulmonary changes	Nishiyama et al., 1975
Monkey	5 months	0.7	pulmonary	pulmonary changes	Nishiyama et al., 1975
Monkey	10 months	0.7	pulmonary	inflammatory and radiographic changes	Suzuki et al., 1977
Mouse	14 days	3.0	pulmonary	reversible inflammatory changes	Nishiyama et al., 1975
Monkey	5 months	3.0	pulmonary	congestion	Nishiyama et al., 1975
Monkey	10 months	3.0	pulmonary	inflammatory and radiographic changes	Suzuki et al., 1978
Guinea pig	1 day	22	pulmonary	inflammatory reaction with bacterial challenge	Bergström, 1977
Mouse	1-4 days	109	pulmonary; mortality rate	increased mortality with bacterial challenge	Malgetter et al., 1976



TABLE 9-1 (cont.)

Species	Duration	Concentration (mg/m <sup>3</sup> )	System Examined	Effects*	Reference
<chem>Mn3O4</chem>					
Monkey	15 months	0.10	pulmonary neurologic	none observed	Coulston and Griffin, 1977
Rat, monkey	9 months	1.1	pulmonary neurologic	none observed	Ulrich, 1979c
Rat, monkey	9 months	0.11	pulmonary neurologic	none observed	Ulrich, 1979c
Rat	2 months	0.12 + engine exhaust	pulmonary	none observed	Moore et al., 1975
Hamster	2 months	0.12	pulmonary	none observed	Moore et al., 1975
Mouse	2 hours	0.89	pulmonary	none observed	Adkins et al., 1980a
Mouse	2 hours	0.22-2.66 + bacteria	mortality rate	increases in rate	Adkins et al., 1980c

\*These results are described in greater detail in Chapter 6 and in Table 6-10.

Table 9-1 summarizes several studies which report no gross or microscopic changes after exposure to  $\sim 0.1 \text{ mg/m}^3 \text{ Mn}_3\text{O}_4$ .

Thus, the animal data qualitatively support a range of respiratory effects associated with exposure to manganese. Human data qualitatively describe such effects but have limited exposure/response information because exposure ranges are broad, cohorts are not followed for long time periods, and duration of exposure is unreported or variable within a study population.

The mechanisms for toxic effects other than carcinogenicity are consistent with the concept of a threshold. The conventional approach toward determining the threshold for noncarcinogenic toxicity is to bracket it by identifying the highest level at which no adverse effects are observed (NOEL) and the lowest level at which adverse effects are observed (LOAEL). Therefore, the health effects assessment for manganese, considering the data available, focuses on the highest NOELs in humans or on the LOAEL as available. These data can be supported by animal data by estimating human equivalent exposures from animal exposure/effect levels.

#### 9.5. HEALTH HAZARD EVALUATION

9.5.1. Critical Effect and Effect Levels. The critical effect is that adverse health effect which occurs at the lowest level of exposure. In order to identify the critical effect, the highest no-observed-effect-level (NOEL) and the lowest-observed-adverse-effect-level (LOAEL) for relevant toxic effects are identified and compared. Qualitative results and dose/response data from experimental animals are compared with levels based on human experience. Studies in humans report effects on the respiratory symptoms at levels below  $1 \text{ mg/m}^3$  whereas studies of effects on the CNS below this level are equivocal or negative. Two studies give exposure-response

information in humans for the clinical field. Nogawa et al. (1973) reported an increased prevalence of respiratory symptoms in schoolchildren exposed to 0.005-0.04  $\text{mg}/\text{m}^3$  manganese emissions from a ferromanganese plant. This is in contrast to the results of Santic and Lucic-Palavic (1977) report an increased prevalence of chronic bronchitis in workers exposed to 0.4-16  $\text{mg}/\text{m}^3$ ; however, prevalence of chronic bronchitis in a group of workers exposed to 0.005-0.04  $\text{mg}/\text{m}^3$  did not differ from controls. These results do not contradict the results of Nogawa because 1) children may be expected to be more sensitive than male workers, and 2) the latter study had less statistical power because fewer subjects were involved.

NOELs could be derived from several studies reported in laboratory animals exposed to manganese oxides consisting largely of particles in the alveolar fraction ( $<2 \mu\text{m}$ , see Section 3.6.4.1.). These studies are summarized in Tables 6-10 and 9-1. Factors which compromise the use of these studies for NOELs are described below.

Coulston and Griffin (1977) did not perform tests of lung function, did not give details of the pathological examination and reported acariasis and associated pulmonary complications in a majority (8/12) of the animals studied. Moore et al. (1975) reported no gross or microscopic abnormalities; however, they observed the animals for only 8 hours/day and for only 56 days. Ulrich et al. (1979a,b,c) exposed rats and squirrel monkeys to three levels of manganese and a control for 9 months. Pulmonary function tests were performed only on the monkeys. This study also had deficiencies which reduced confidence in the range of negative findings reported. Due to the small group size of the monkeys (4 males and 4 females) and the large within group variability, it lacked sufficient statistical power to detect any but the largest changes in the parameters measured. The variability

could have been reduced by using more appropriate statistical analysis to control for within group variation. The description of lung pathology was inadequate. Negative results were reported at  $1.15 \text{ mg/m}^3$ ; however, Suzuki et al. (1978) reported pathologic changes in the lungs of rhesus monkeys exposed to  $700 \text{ } \mu\text{g/m}^3$  of  $\text{MnO}_2$  for 10 months. Based on these data, the next highest NOEL reported by Ulrich et al. (1979a,b,c) was  $0.113 \text{ mg/m}^3$ . However, the repeated reports of the absence of gross and microscopic abnormalities at a similar level ( $0.1 \text{ mg/m}^3$ ) suggest that this level may be close to a threshold.

These data do not exclude the possibility that more subtle toxic effects on the lungs may occur at  $\sim 0.1 \text{ mg/m}^3$ . Effects do occur at  $0.7 \text{ mg/m}^3$ . In order to compare the reported NOEL ( $0.1 \text{ mg/m}^3$ ) and the LOAEL ( $0.7 \text{ mg/m}^3$ ) to similar data in humans, it would be helpful to estimate a human intake equivalent to that of the experimental animals. The suggested approach is provided in the Appendix.

## 10. REFERENCES

Abernathy, R.F., M.J. Peterson and F.H. Gibson. 1969. Spectrochemical analysis of coal ash for trace elements. Bur. Mines Report of Investigations 7281. U.S. Dept. Interior, Pittsburgh, PA. 30 p.

Abrams, E., J.W. Lassiter, W.J. Miller, M.W. Neathery and R.P. Gentry. 1976a. Effect of dietary manganese as a factor affecting  $^{54}\text{Mn}$  absorption in rats. Nutr. Rep. Inter. 14(5): 561-565.

Abrams, E., J.W. Lassiter, W.J. Miller, M.W. Neathery, R.P. Gentry and R.D. Scarth. 1976b. Absorption as a factor in manganese homeostasis. J. Anim. Sci. 42(3): 630-636.

ACGIH (American Conference of Governmental Industrial Hygienists). 1958. Committee on Recommended Analytical Methods: Determination of manganese in air, periodate oxidation method. In: Manual of Analytical Methods Recommended for Sampling and Analysis of Atmospheric Contaminants. ACGIH, Cincinnati, OH. Mn-1, Mn-4.

ACGIH (American Conference of Governmental and Industrial Hygienists). 1980. Documentation of the Threshold Limit Values, fourth ed. ACGIH, Cincinnati, OH. p. 250-251.

Ad Hoc Working Group of Technical Committee - Air Quality, International Standards Organization. 1981. Recommendations on size definitions for particle sampling. Am. Ind. Hyg. Assoc. J. 42: A64-A68.

Adkins, B., Jr., G.H. Luginbuhl and D.E. Gardner. 1980a. Acute exposure of laboratory mice to manganese oxide. *Am. Ind. Hyg. Assoc. J.* 41: 494-500.

Adkins, B., Jr., G.H. Luginbuhl and D.E. Gardner. 1980b. Biochemical changes in pulmonary cells following manganese oxide inhalation. *J. Toxicol. Environ. Health.* 6: 445-454.

Adkins, B., Jr., G.H. Luginbuhl, F.J. Miller and D.E. Gardner. 1980c. Increased pulmonary susceptibility to streptococcal infection following inhalation of manganese oxide. *Environ. Res.* 23: 110-120.

Ahrland, D.S. 1975. Metal complexes present in seawater. In: *The Nature of Seawater*, D. Goldberg, Ed. Dahlem, Kopenhagen, Berlin. p. 219-244.

Ajemian, R.S. and N.E. Whitman. 1969. Determination of manganese in urine by atomic absorption spectrometry. *Am. Ind. Hyg. Assoc. J.* 30: 52-56.

Alexander, F.W., B.E. Clayton and H.T. Delves. 1974. Mineral and trace-metal balances in children receiving normal and synthetic diets. *Q. J. Med.* 43: 89-111.

Alexander, M. 1967. *Introduction to Soil Microbiology*. John Wiley and Sons, Inc., New York. 472 p.

Alexander, M. 1977. *Introduction to Soil Microbiology*, 2nd ed. John Wiley and Sons, Inc., New York. p. 385-392.

Alpert, D.J. and P.K. Hopke. 1981. A determination of the sources of airborne particles collected during the Regional Air Pollution Study. *Atmos. Environ.* 15: 675-687.

Amdur, Mo., L.C. Norris and G.F. Heuser. 1945. The need for manganese in bone development in the rat. *Proc. Soc. Exp. Biol. Med.* 59: 254-255.

Anke, M. and H.J. Schneider. 1974. Trace element concentrations in human kidney in relation to age and sex. *Zschr. Urol.* 67: 357-362.

Ansola, J., E. Uiberall and E. Escudero. 1944a. Intoxication by manganese in Chile (study on 64 cases). I. Environmental and etiological factors. *Rev. Med. Chile.* 72: 222-228.

Ansola, J., E. Uiberall and E. Escudero. 1944b. Intoxication by manganese in Chile (study on 64 cases). II. Clinical aspects, incapacity and medico-legal reparations. *Rev. Med. Chile.* 72: 311-322.

Ardid, R.R. and A.S. Torrente. 1949. A case of amylostatic syndrome due to occupational poisoning with manganese. *Actas Espan. Neurol. Psiquiat.* 8: 42-54.

Ashizawa, R. 1927. Report on an autopsy case of chronic manganese poisoning. *Jap. J. Med. Sci. Trans.* 7: 173-191.

Autissier, N., L. Rochette, P. Dumas, A. Belay, A. Loireau and J. Bralet. 1982. Dopamine and norepinephrine turnover in various regions of the rat brain after chronic manganese chloride administration. *Toxicology*. 24: 175-182.

Baader, E.W. 1932. Manganvergiftungen in Elementefabriken. (Manganese poisoning in dry cell battery factories). *Arch. Gewerbepath. Gewerbehyg.* 4: 101-116. (Ger.)

Baader, E.W. 1937. Die Manganpneumonie. *Arztl. Sachverst. Ztg.* 43: 75-81. (Ger.)

Baader, E.W. 1939. Eine Restudie über schweren Manganismus bei ägyptischen Mangangrubenarbeitern. *Arch. Gewerbepath. Gewerbehyg.* 9: 477-486. (Ger.)

Bacon, F.E. 1967. Manganese and manganese alloys. In: Kirk-Othmer Encyclopedia of Chemical Technology, Second ed., Vol. 12, H.F. Mark et al., Ed. John Wiley and Sons, Inc., New York. p. 887-905.

Baetz, R.A. and C.T. Kenner. 1975. Determination of trace metals in foods using chelating ion exchange concentration. *Agric. Food Chem.* 23: 41-45.

Balaní, S.G., G.M. Umarji, R.A. Bellare and H.C. Merchant. 1967. Chronic manganese poisoning. A case report. *J. Postgrad. Med.* 13: 116-122.

Barbeau, A., N. Inou'e and T. Cloutier. 1976. Role of manganese in dystonia. *Adv. Neurol.* 14: 339-352.



Barrows, E.H. 1983. Filter Bank data on Manganese, 1977-1982. Letter from E.H. Barrows, Northrop Services, Inc., Research Triangle Park, NC to R.J.F. Bruins, Environmental Criteria and Assessment Office, U.S. Environmental Protection Agency, Cincinnati, OH. April 22.

Baxter, D.J., W.O. Smith and G.C. Klein. 1965. Some effects of acute manganese excess in rats. Proc. Soc. Exp. Biol. Med. 119: 966-970.

Bear, F.E. 1964. Chemistry of the Soil. Reinhold Publishing Co., New York. p. 342.

Belz, R. 1960. The amounts of iron, copper, manganese and cobalt in average diets of various age groups in the Netherlands. Voeding. 21: 236-251.

Bergström, R. 1977. Acute pulmonary toxicity of manganese dioxide. Scand. J. Work Environ. Health. 3: 1-40.

Bernheimer, H., W. Birkmayer, O. Hornykiewicz, K. Jellinger and F. Seitelberger. 1973. Brain dopamine and the syndromes of Parkinson and Huntington-Clinical, morphological and neurochemical correlations. J. Neurol. Sci. 20: 415-425.

Bernstein, D.M. and K.A. Rahn. 1979. New York summer aerosol study: Trace element concentration as a function of particle size. Ann. N.Y. Acad. Sci. 322: 87-97.

Bernstein, D.M., M.T. Kleinman, T.J. Kneip, T.L. Chan and M. Lippmann. 1976. A high-volume sampler for the determination of particle size distributions in ambient air. J. Air Pollut. Control Assoc. 26: 1069-1072.

Bertinchamps, A.J. and G.C. Cotzias. 1958. Biliary excretion of manganese. Fed. Proc. 17: 428.

Bertinchamps, A.J., S.T. Miller and G.C. Cotzias. 1966. Interdependence of routes excreting manganese. Am. J. Physiol. 211(1): 217-224.

Bethard, W.F., D.A. Olehy and R.A. Schmitt. 1964. The use of neutron activation analysis for the quantitation of selected cations in human blood. In: L'analyse par Radioactivation et les Applications aux Sciences Biologiques. III. Colloque International de Biologie de Saclay, Sept. 1963. Presses Univ. France, Paris. p. 379-393.

Blenstock, D. and F.J. Field. 1960. Bench-scale investigation on removing sulfur dioxide from flue gases. J. Air Pollut. Control Admin. 10: 121-125.

Bienvenu, P., C. Noire and A. Cler. 1963. Comparative general toxicity of metallic ions. A relation with the periodic classification. Rech. Serv. Sante Armees, Lyons, France. 256: 1043-1044.

Bize, I.B. and L.W. Oberly. 1979. Superoxide radical production and superoxide dismutase content of hepatoma mitochondria and normal liver. Proc. Am. Assoc. Cancer Res. 20: 139. (Abstr.)

Bize, I.B., L.W. Oberly and H.P. Morris. 1980. Superoxide dismutase and superoxide radical in Morris hepatomas. *Cancer Res.* 40: 3686-3693.

Bond, A.M. and B.M. Kelly. 1977. Investigation of automated differential pulse anodic stripping voltammetry of the problem of storage of dilute solutions. *Atlanta.* 24: 453-457.

Bonilla, E. 1978a. Flameless atomic absorption spectrophotometric determination of manganese in rat brain and other tissues. *Clin. Chem.* 24(3): 471-474.

Bonilla, E. 1978b. Increased GABA content in caudate nucleus of rats after chronic manganese chloride administration. *J. Neurochem.* 31: 551-552.

Bonilla, E. 1980. L-tyrosine hydroxylase activity in the rat brain after chronic oral administration of manganese chloride. *Neurobehav. Toxicol.* 2: 37-41.

Bonilla, E. 1983. Chronic manganese poisoning and striatal adenylate cyclase activity. *Trans. Am. Soc. Neurochem.* 14: 123.

Bonilla, E. and M. Diez-Ewald. 1974. Effect of L-dopa on brain concentration of dopamine and homovanillic acid in rats after chronic manganese chloride administration. *J. Neurochem.* 22: 297-299.

Bonilla, E. and H. Martinez. 1981. The effects of manganese on cholinergic activity in rat brain. *Trans. Am. Soc. Neurochem.* 12: 246.

Boutron, C. and S. Martin. 1979. Preconcentration of dilute solutions at the  $10^{-12}$  g/g level by nonboiling evaporation with variable variance calibration curves. Anal. Chem. 51: 140-145.

Boyce, W. and C.L. Witzleben. 1973. Bilirubin as a cholestatic agent. II. Effect of variable doses of bilirubin on the severity of manganese-bilirubin cholestasis. Am. J. Pathol. 72: 427-432.

Bracewell, J.M. and D. Call. 1967. The catalytic oxidation of sulfur dioxide in solution at concentrations occurring in fog droplets. In: Air Pollution, Proceedings of the Symposium on the Physico-Chemical Transformation of Sulphur Compounds in the Atmosphere and the Formation of Acid Smogs, Mainz, Germany, June 8-9, 1967. Organization for Economic Cooperation and Development, Directorate for Scientific Affairs. (Cited in Sullivan, 1969)

Breck, W.G. 1974. Redox levels in the sea. In: The Sea, Vol. 5, E.D. Goldberg, Ed. Marine Chemistry. Wiley-Interscience, New York.

Brozina, E. 1921. Internationale Übersicht über Gewerbekrankheiten nach den Berichten der Gewerbeinspektionen der K. K. Länder über das Jahr 1913. J. Springer, Berlin. 8: 143. (Ger.)

Britton, A.A. and G.C. Cotzias. 1966. Dependence of manganese turnover on intake. Am. J. Physiol. 211(1): 203-206.

Bubarev, A. 1931. On the poisoning associated with loading and unloading manganese on ships. Gig. Bezopasn. Pat. Truda. 9: 85-87.

Buchet, J.P., R. Lauwerys, H. Roels and C. de Vos. 1976. Determination of manganese in blood and urine by flameless atomic absorption spectrophotometry. Clin. Chim. Acta. 73: 481-486.

Buono, J.A., R.W. Karin and J.L. Fasching. 1975. The rapid separation of trace metals from neutron-activated saline matrices by chelation on a poly-5-vinyl-8-hydroxyquinoline column. Anal. Chim. Acta. 80: 327-334.

Burch, R.E., R.V. Williams, H.K.J. Hahn, M.M. Jetton and J.F. Sullivan. 1975. Tissue trace element and enzyme content in pigs fed a low manganese diet. I. A relationship between manganese and selenium. J. Lab. Clin. Med. 86: 132-139.

Burnett, W.T., Jr., R.R. Bigelow, A.W. Kimball and C.W. Sheppard. 1952. Radio-manganese studies on the mouse, rat and pancreatic fistula dog. Am. J. Physiol. 168: 620-625.

Büttner, H.E. and E. Lenz. 1937. Über die Möglichkeit von Manganschaden im Braunkohlenbergwerk. Arch. Gewerbepath. Gewerbehyg. 7: 672-684. (Ger.)

Cahill, D.F., M.S. Bercegeay, R.C. Haggerty, J.E. Gerding and L.E. Gray. 1980. Age-related retention and distribution of ingested  $Mn_3O_4$  in the rat. Toxicol. Appl. Pharmacol. 53: 85-91.

Calabrese, E.J. 1983. Principles of Animal Extrapolation. John Wiley and Sons, New York.

Canavan, M.M., S. Cobb and C.K. Drinker. 1934. Chronic manganese poisoning. Report of a case, with autopsy. Arch. Neurol. Psychiatr. 32: 501-513.

Carter, S.D., J.F. Hein, G.L. Rehnberg and J.W. Laskey. 1980. Chronic manganese oxide ingestion in rats: Hematological effects. J. Toxicol. Environ. Health. 6: 207-216.

Casamajor, L. 1913. An unusual form of mineral poisoning affecting the nervous system: Manganese. J. Am. Med. Assoc. 69: 646-649.

Casey, C.E. and M.F. Robinson. 1978. Copper, manganese, zinc, nickel, cadmium and lead in human foetal tissues. Br. J. Nutr. 39: 639-646.

Cauvin, F. 1943. Les maladies professionnelles des mineurs du manganese. Maroc. Med. 22: 16. (Cited by Rodier, 1955) (Fre.)

Chan, A.W.K., L.K.J. Lai, M. Minski, L. Lim and A.N. Davison. 1981. Manganese concentration in rat organs: Effect after life-long manganese treatment. Biochem. Soc. Trans. 9: 229.

Chan, A.W.K., M.J. Minski, L. Lim and J.C.K. Lai. 1983. Changes in brain regional manganese levels in post-natal development: Modulations by chronic manganese administration. Neurotoxicology (Mn Symposium Issue) (In press)

Chandra, S.V. 1971. Cellular changes induced by manganese in the rat testis. Acta Pharmacol. Toxicol. 29: 75-80.

Chandra, S.V. 1972. Histological and histochemical changes in experimental manganese encephalopathy in rabbits. Arch. Toxicol. 29: 29-38.

Chandra, S.V. and Z. Imam. 1973. Manganese induced histochemical and histological alterations in gastrointestinal mucosa of guinea pig. Acta Pharmacol. Toxicol. 33: 449-458.

Chandra, S.V. and Z. Imam. 1975. Effect of manganese on the morphology of the rabbit adrenal cortex. Arch. Hig. Rada. 26: 201-208.

Chandra, S.V. and G.S. Shukla. 1978. Manganese encephalopathy in growing rats. Environ. Res. 15: 28-37.

Chandra, S.V. and G.S. Shukla. 1981. Concentrations of striatal catecholamines in rats given manganese chloride through drinking water. J. Neurochem. 36(2): 683-687.

Chandra, S.V. and S.P. Srivastava. 1970. Experimental production of early brain lesions in rats by parenteral administration of manganese chloride. Acta Pharmacol. Toxicol. 28: 177-183.

Chandra, S.V. and S.K. Tandon. 1973. Enhanced manganese toxicity in iron-deficient rats. Environ. Physiol. Biochem. 3: 230-235.

Chandra, S.V., R. Ara, M. Nagar and P.K. Seth. 1975a. Sterility in experimental manganese toxicity. Acta Biol. Med. Ger. 30: 857-862.

Chandra, S.V., Z. Imam and N. Nagar. 1973b. Significance of serum calcium, inorganic phosphates and alkaline phosphatase in experimental manganese toxicity. *Ind. Health.* 11: 43-47.

Chandra, S.V., P.K. Seth and J.K. Mankeshwar. 1974. Manganese poisoning: Clinical and biochemical observations. *Environ. Res.* 7: 374-380.

Chandra, S.V., D.K. Saxena and M.Z. Hasan. 1975. Effect of zinc on manganese induced testicular injury in rats. *Ind. Health.* 13: 51-56.

Chandra, S.V., G.S. Shukla and D.K. Saxena. 1979a. Manganese-induced behavioral dysfunction and its neurochemical mechanism in growing mice. *J. Neurochem.* 33: 1217-1221.

Chandra, S.V., G.S. Shukla and R.C. Murthy. 1979b. Effect of stress on the response of rat brain to manganese. *Toxicol. Appl. Pharmacol.* 47: 603-608.

Chandra, S.V., R.S. Srivastava and G.S. Shukla. 1979c. Regional distribution of metals and biogenic amines in the brain of monkeys exposed to manganese. *Toxicol. Lett.* 4: 189-192.

Chandra, S.V., G.S. Shukla, R.S. Srivastava, H. Singh and V.P. Gupta. 1981. An exploratory study of manganese exposure to welders. *Clin. Toxicol.* 18: 407-416.

Chittleborough, G. 1980. A chemist's view of the analysis of human hair for trace elements. *Sci. Total Environ.* 14: 53-75.



Cikrt, M. 1972. Biliary excretion of  $^{203}\text{Hg}$ ,  $^{64}\text{Cu}$ ,  $^{52}\text{Mn}$ , and  $^{210}\text{Pb}$  in the rat. Br. J. Ind. Med. 29: 74-80.

Cikrt, M. 1973. Enterohepatic circulation of  $^{64}\text{Cu}$ ,  $^{52}\text{Mn}$  and  $^{203}\text{Hg}$  in rats. Arch. Toxicol. 34: 51-59.

Cikrt, M. and J. Vostal. 1969. Study of manganese resorption in vitro through intestinal wall. Int. J. Clin. Pharmacol. 2: 280-285.

Coe, M., R. Cruz and J.C. van Loon. 1980. Determination of methylcyclopentadienylmanganesetricarbonyl by gas chromatography-atomic absorption spectrometry at  $\text{ng m}^{-3}$  levels in air samples. Anal. Chim. Acta. 120: 171-176.

Cook, D.G., S. Fahn and K.A. Bratt. 1974. Chronic manganese intoxication. Arch. Neurol. 30: 59-64.

Cooper, J., F. Bloom and R. Roth. 1982. The Biochemical Basis of Neuropharmacology, 4th ed. Oxford University Press, New York.

Cooper, J.A. and J.G. Watson, Jr. 1980. Receptor oriented methods of air particulate source apportionment. J. Air Pollut. Control Assoc. 30: 1116-1125.

Cotzias, G.C. 1958. Manganese in health and disease. Physiol. Revs. 38: 503-533.

Cotzias, G.C. 1962. Manganese. In: Mineral Metabolism: An Advanced Treatise, 28, C.L. Comar and F. Bronner, Ed. Academic Press, New York. p. 403-442.

Cotzias, G.C. and P.S. Papavasiliou. 1962. State of binding of natural manganese in human cerebrospinal fluid, blood and plasma. *Nature* (London). 195: 823-834.

Cotzias, G.C., S.T. Miller and J. Edwards. 1966. Neutron activation analysis: The stability of manganese concentrations in human blood and serum. *J. Lab. Clin. Med.* 67: 836-849.

Cotzias, G.C., K. Horiuchi, S. Fuenzalida and I. Mena. 1968. Chronic manganese poisoning. Clearance of tissue manganese concentrations with persistence of the neurological picture. *Neurology*. 18: 376-382.

Cotzias, G.C., P.S. Papavasiliou, J. Ginos, A. Steck and S. Oüby. 1971. Metabolic modification of Parkinson's disease and of chronic manganese poisoning. *Ann. Rev. Med.* 22: 305-326.

Cotzias, G.C., S.T. Miller, P.S. Papavasiliou and L.C. Tang. 1976. Interactions between manganese and brain dopamine. *Med. Clin. North Am.* 60: 729-738.

Coulston, F. and T. Griffin. 1977. Inhalation toxicology of airborne particulate manganese in rhesus monkeys. EPA 600/1-77-026. PB 268 643.

Crowther, J. 1978. Semiautomated procedure for the determination of low levels of total manganese. *Anal. Chem.* 50: 1041-1043.

Cullimore, D.R. and A.E. McCann. 1977. The identification, cultivation and control of iron bacteria in groundwater. In: *Aquatic Microbiology*, F.A. Skinner and J.M. Shewan, Ed. Academic Press, London. p. 219-261.

Curzon, G. 1975. Metals and melanins in the extrapyramidal centers. *Pharmacol. Ther.* 8. 1: 673-684.

Dagli, A.J., D. Golden, M. Finkel and E. Austin. 1973. Pyloric stenosis following ingestion of potassium permanganate. *Digestive Diseases.* 18: 1091-1094.

D'Amico, D.J. and H.L. Klawans. 1976. Direct microdetermination of manganese in normal serum and cerebrospinal fluid by flameless atomic absorption spectrophotometry. *Anal. Chem.* 48: 1469-1472.

Daniels, A.J., K. Gysling and J. Abarca. 1981. Uptake and release of manganese by rat striatal slices. *Biochem. Pharmacol.* 30(13): 1833-1837.

Dantin Gallego, J. 1935. L'Hygiene et la pathologie dans le travail avec le manganese. Instituto Nacional de Prevision Publ. 445, Madrid. (Fre.)

Dantin Gallego, J. 1944. Intoxicacion por el manganese. In: *Curso de Higiene del Trabajo*. Publicaciones de la Jefatura Provincial de Sanidad, Madrid. p. 459-473. (Spa.)

Dastur, D.K., D.K. Manghani, K.V. Raghavendran and K.M. Jeejeebhoy. 1969. Distribution and fate of  $Mn^{54}$  in the rat, with special reference to the CNS. Q. J. Exp. Physiol. 54: 322-331.

Dastur, D.K., D.K. Manghani and K.V. Raghavendran. 1971. Distribution and fate of  $Mn^{54}$  in the monkey: Studies of different parts of the central nervous system and other organs. J. Clin. Invest. 50: 9-20.

Davies, C.N., M. Aylward and D. Leacey. 1951. Impingement of dust from air jets. Arch. Ind. Hyg. Occup. Health. 4: 354-397.

Davis, B.L., L.R. Johnson, R.K. Stevens, W.J. Courtney and D.W. Safriet. 1984. The quartz and elemental composition of aerosols from 22 U.S. cities of the EPA inhalable particulate network. Atmos. Environ. 18: In press.

Davison, R.L., D.F.S. Natusch, J.R. Wallace and C.A. Evans, Jr. 1974. Trace elements in fly ash. Dependence of concentration on particle size. Environ. Sci. Technol. 8: 1107-1113.

DeHuff, G.L. 1961-1980. Manganese. In: Minerals Yearbooks, 1960-1979. Vol. 1. U.S. Bureau of Mines, Dept. of Interior, Washington, DC.

DeHuff, G.L. 1973. Manganese in the ecosystem. In: Medical and Biologic Effects of Environmental Pollutants: Manganese. NRC, NAS, Washington, DC. p. 3-50.

DeHuff, G.L. 1975. Manganese. In: Mineral Facts and Problems. Bur. Mines Bull. 667. U.S. Dept. Interior, Washington, DC. p. 653-668.

DeHuff, G.L. and T.S. Jones. 1980. Manganese Preprint from Bulletin 671 -- Mineral Facts and Problems, 1980 ed. Bur. Mines, U.S. Dept. Interior, Washington, DC.

DeHuff, G.L. and T.S. Jones. 1981. Manganese. In: Minerals Yearbook 1980. Vol. 1. Metals and Minerals. U.S. Bureau of Mines, U.S. Dept. of Interior, Washington, DC.

De Lamirande, E. and G.L. Plaa. 1978. Role of manganese, bilirubin and sulfobromophthalein in manganese-bilirubin cholestasis in rats. Proc. Soc. Exp. Biol. Med. 158: 283-287.

De Lamirande, E. and G.L. Plaa. 1979a. Dose and time relationships in manganese-bilirubin cholestasis. Toxicol. Appl. Pharmacol. 49: 257-263.

De Lamirande, E. and G.L. Plaa. 1979b. Bilirubin excretion pattern in manganese-bilirubin cholestasis. Arch. Int. Pharmacol. Therap. 239: 24-35.

Delves, H.T. 1970. A micro-sampling method for the rapid determination of lead in blood by atomic-absorption spectrophotometry. Anal. 95: 431-438.

Deskin, R., S.J. Bursian and F.W. Edens. 1981a. Neurochemical alterations induced by manganese chloride in neonatal rats. Neurotoxicology. 2: 65-73.

Deskin, R., S.J. Bursian and F.W. Edens. 1981b. An investigation into the effects of manganese and other divalent cations on tyrosine hydroxylase activity. *Neurotoxicology*. 2: 75-81.

Diez-Ewald, M., L.R. Weintraub and W.H. Crosby. 1968. Interrelationship of iron and manganese metabolism. *Proc. Soc. Exp. Biol. Med.* 129: 448-451.

Dikshith, T.S. and S.V. Chandra. 1978. Cytological studies in albino rats after oral administration of manganese chloride. *Bull. Environ. Contam. Toxicol.* 19: 741-746.

DiPaolo, J.A. 1964. The potentiation of lymphosarcomas in mice by manganese chloride. *Fed. Proc.* 23: 393.

Doi, M. 1959. Experimental studies on the chronic manganese poisoning. *Shikoku Igaku Zasshi*. 15: 1789-1802.

Dokuchaev, V.F. and N.N. Skvortsova. 1962. Atmospheric air pollution with manganese compounds and their effect on the organism. In: *USSR Literature on Air Pollution and Related Occupational Diseases, Part 1*. Meditsina Press. 9: 40-46.

Donaldson, J. 1981. The pathophysiology of trace metal: Neurotransmitter interaction in the CNS. *Trends Pharmacol. Sci.* 2: 75-78.

Donaldson, J., F.S. LaBella and D. Gesser. 1981. Enhanced autoxidation of dopamine as a possible basis of manganese neurotoxicity. *Neurotoxicology*. 2: 53-64.

Donaldson, J., J. McGregor and F. LaBella. 1982. Manganese neurotoxicity: A model for free radical mediated neurodegeneration? Can. J. Physiol. Pharmacol. 60: 1398-1405.

Dourson, M.L. and J.F. Stara. 1983. Regulatory history and experimental support of uncertainty (safety) factors. Reg. Toxicol. Pharmacol. (In press)

Doyle, J.J. and W.H. Pfander. 1975. Interactions of radium with copper, iron, zinc, and manganese in ovine tissues. J. Nutr. 105: 599-606.

Durfor, C.N. and E. Becker. 1964. Public water supplies of the 100 largest cities in the United States. U.S. Geol. Sur., Water-Supply Paper 1812, Washington, DC. 364 p.

Durum, W.H. and J. Haffty. 1963. Implications of the minor element content of some major streams of the world. Geochim. Cosmochim. Acta. 27: 1-11.

Dzubay, T.G. 1980. Chemical elemental balance method applied to dichotomous sampler data. Ann. New York Acad. Sci. 338: 126-144.

Dzubay, T.G. and R.O. Nelson. 1975. Self absorption corrections for X-ray fluorescence analysis of aerosols. In: Advances in X-Ray Analysis, Vol. 18, W.L. Pickles, C.S. Barrett, J.B. Newkirk and C.O. Ruud, Ed. p. 619-631.

Dzubay, T.G. and R.K. Stevens. 1975. Ambient air analysis with dichotomous sampler and X-ray fluorescence spectrometer. Environ. Sci. Technol. 9: 663-668.

Dzubay, T.G., R.K. Stevens, W.J. Courtney and E.A. Drane. 1981. Chemical element balance analysis of Denver aerosol. In: Electron Microscopy and X-Ray Applications to Environmental and Occupational Health Analysis, Vol 2, P.A. Russell, Ed. Ann Arbor Science Publ. Inc. p. 23-42.

Dzubay, T.G., R.K. Stevens, C.W. Lewis, et al. 1982. Visibility and aerosol composition in Houston, TX. Environ. Sci. Technol. 16: 514-525.

Eads, E.A. and C.E. Lambdin. 1973. Survey of trace metals in human hair. Environ. Res. 6: 247-252.

Ehrlich, H.L. 1972. Response of some activities of ferromanganese nodule bacteria to hydrostatic pressure. In: Effect of the Ocean Environment on Microbial Activities, R.R. Colwell and R.Y. Morita, Ed. University Park Press, Baltimore. p. 208-221.

Ellis, G.H., S.E. Smith and E.M. Gates. 1947. Further studies of manganese deficiency in the rabbit. J. Nutr. 34: 21-31.

Elstad, O. 1939. Factory smoke containing manganese as predisposing cause in epidemics of pneumonia in an industrial district. Nord. Med. 3: 2527-2532.

Emara, A.M., S.H. El-Ghawabi, O.I. Madkour and G.H. El-Sarma. 1971. Chronic manganese poisoning in the dry battery industry. Br. J. Ind. Med. 28: 73-82.



Energy Information Administration. 1980. Annual Report to Congress. Vol. 2. p. 7, 63.

Epstein, S S., E. Arnold, J. Andra, W. Bass and Y. Bishop. 1972. Detection of chemical mutagens by the dominant lethal assay in the mouse. Toxicol. Appl. Pharmacol. 23: 288-325.

Erickson, C.K. 1978. Functional relationships among central neurotransmitters. Rev. Neurosci. 3: 1-34.

Ethyl Corporation. 1972. Information for the National Research Council concerning methylcyclopentadienyl manganese tricarbonyl. Ethyl Corp. Res. Labs., Detroit, MI. September 8.

Ethyl Corporation. 1973. The effect of manganese on the oxidation of sulfur dioxide in the air. Ethyl Corp. Res. Labs., Detroit, MI. November 28.

Everson, G.J., L.S. Hurley and J.F. Geiger. 1959. Manganese deficiency in the guinea pig. J. Nutr. 68: 49.

Fairhall, L.T. and P.A. Neal. 1943. Industrial manganese poisoning. Natl. Inst. Health Bull. No. 182. Washington, DC. p. 24.

FAO/WHO (Food Agricultural Organization/World Health Organization). 1977. Report of the Consultation on the Joint FAO/WHO Food and Animal Feed Contamination Monitoring Programme - Phase II, Geneva.

Fee, J.A. 1980. Superoxide, superoxide dimatases and oxygen toxicity. In: Metal Ion Activation of Dioxygen, Vol. 2 of Metal Ions in Biology, T.G. Spiro, Ed. Wiley-Interscience, New York. p. 209-237.

Fischer, H. and P. Weigert. 1977. Die Schwermetalle Kupfer, Quecksilber, Mangan und Zink in menschlichen Organen. Jff. Gesundh. Wesen. 39: 269-278. (Ger.)

Flato, J.B. 1972. The renaissance in polarographic and voltammetric analysis. Anal. Chem. 44: 75A-87A.

Flinn, R.H., P.A. Neal, W.A. Reinhart, J.M. Dallavalle, W.B. Fulton and A.E. Dooley. 1940. Chronic manganese poisoning in an ore-crushing mill. Pub. Health Bull. No. 247. Washington, DC. p. 1-77.

Flinn, R.H., P.A. Neal and W.B. Fulton. 1941. Industrial manganese poisoning. J. Ind. Hyg. Toxicol. 23: 374-387.

Fore, H. and R.A. Morton. 1952. Manganese in rabbit tissues. Biochem. J. 51: 600-603.

Franz, R.D. 1962. Toxicities of some trace metals. Naunyn-Schmiedeberg's Arch. Exp. Path. Pharmac. 244: 17-20.

Freise, F.W. 1933. Gesundheitsschadigungen durch Verarbeitung pflanzlicher Drogen. Beobachtungen aus brasilianischen Gewerbebetrieben. Arch. Gewerbe-path. Gewerbehyg. 4: 381-399. (Ger.)

Fugas, M. 1980. Personal communication.

Fukai, R. and Huynh-Ngoc. 1968. Studies on the chemical behavior of radio-nuclides in seawater. I. General considerations and study of precipitation of trace amounts of chromium, manganese, iron, cobalt, zinc and cerium. IAEA Publ. 22. 26 p.

Furst, A. 1978. Tumorigenic effect of an organo-manganese compound on F344 rats and Swiss albino mice. J. Natl. Cancer Inst. 60: 1171.

Gale, N.L., B.G. Wixson, M.G. Hardie and J.C. Jennett. 1973. Aquatic organisms and heavy metals in Missouri's new lead belt. Water Res. Bull. 9: 673-688.

Garcia Avila, M. and R. Penalver. 1953. Manganese poisoning in the mines of Cuba. A preliminary report. Ind. Med. Surg. 22: 220-221.

Garrels, R.M., F.T. Mackenzie and C. Hunt, Ed. 1975. Chemical Cycles and the Global Environment. William Kaufmann, Inc., Los Altos, CA. p. 132.

Gianutsos, G. and M.T. Murray. 1982. Alterations in brain dopamine and GABA following inorganic or organic manganese administration. Neurotoxicology. 3: 75-82.

Gibbons, R.A., S.N. Dixon, K. Hallis, A.M. Russell, B.F. Sansom and H.W. Symonds. 1976. Manganese metabolism in cows and goats. Biochim. Biophys. Acta. 444: 1-10.

Gibson, R.S. and M.S. DeWolfe. 1979. The zinc, copper, manganese, vanadium, and iodine content of hair from 38 Canadian neonates. *Pediat. Res.* 13: 959-962.

Gilfrich, J.V., P.G. Burkhalter and L.S. Birks. 1973. X-ray spectrometry for particulate air pollution -- A quantitative comparison of techniques. *Anal. Chem.* 45: 2002-2009.

Graham, D.G. 1978. Oxidative pathways for catecholamines in the genesis of neuromelanin and cytotoxic quinones. *Mol. Pharmacol.* 14: 633-643.

Graham, D.G. 1983. Catecholamine toxicity: A proposal for the molecular pathogenesis of manganese neurotoxicity and Parkinson's disease. *Neurotoxicology (Mn Special Issue)* (In press)

Graham, D.G., S.M. Tiffany, W.R. Bell, Jr. and W.F. Gutknecht. 1978. Autoxidation versus covalent binding of quinones as the mechanism of toxicity of dopamine, 6-hydroxy-dopamine, and related compounds toward C1300 neuroblastoma cells in vitro. *Mol. Pharmacol.* 14: 644-653.

Graham, J.A., D.E. Gardner, M.D. Waters and D.L. Coffin. 1975. Effect of trace metals on phagocytosis by alveolar macrophages. *Infect. Immun.* 11: 1278-1283.

Gray, L.E., Jr. and J.W. Laskey. 1980. Multivariate analysis of the effects of manganese on the reproductive physiology and behavior of the male house mouse. *J. Toxicol. Environ. Health.* 6: 861-867.

Greathouse, D.G. 1983. Memo from D.G. Greathouse, Health Effects Research Laboratory to R.J.F. Bruins, Environmental Criteria and Assessment Office, U.S. EPA, Cincinnati, OH. March 24.

Greenberg, D.M. and W.W. Campbell. 1940. Studies in mineral metabolism with the aid of induced radioactive isotopes. IV. Manganese. Proc. Natl. Acad. Sci. 26: 448-452.

Greenberg, D.M., D.H. Copp and E.M. Cuthbertson. 1943. Studies in mineral metabolism with the aid of artificial radioactive isotopes. VII. The distribution and excretion, particularly by way of the bile, of iron, cobalt, and manganese. J. Biol. Chem. 147: 749-756.

Gregory, E. and J.T. Staley. 1982. Widespread distribution of ability to oxidize manganese among freshwater bacteria. Appl. Environ. Microbiol. 44: 509-511.

Guthrie, B.E. 1975. Chromium, manganese, copper, zinc and cadmium content of New Zealand foods. N.Z. Med. J. 82: 418-424.

Habibi, K. 1970. Characterization of particulate lead in vehicle exhaust -- environmental techniques. Environ. Sci. Technol. 4: 239-248.

Hagenfeldt, K., L.O. Plantin and E. Diczfalusy. 1973. Trace elements in the human endometrium. II. Zinc, copper, and manganese levels in the endometrium, cervical mucus and plasma. Acta Endocrinol. 72: 115-126.

Hahn, K.J., D.J. Tuma and J.L. Sullivan. 1968. Rapid and simple continuous radiochemical separation of copper, magnesium, zinc, and manganese in biological materials. Anal. Chem. 40: 974-976

Hall, C.A. 1983a. Letter from C.A. Hall, Ethyl Corporation, Ferndale, MI to L.S. Erdreich, Environmental Criteria and Assessment Office, U.S. EPA, Cincinnati, OH. February 18.

Hall, C.A. 1983b. Letter from C.A. Hall, Ethyl Corporation, Ferndale, MI to L.S. Erdreich, Environmental Criteria and Assessment Office, U.S. EPA, Cincinnati, OH. August 1.

Hartman, R.H., G. Matrone and G.H. Wise. 1955. Effect of high dietary manganese on hemoglobin formation. J. Nutr. 57: 429-439.

Hay, J.O. 1967. Manganese compounds. In: Kirk-Othmer Encyclopedia of Chemical Technology, Second ed., Vol. 13, H.F. Mark et al., Ed. Interscience Publishers, New York. p. 1-55.

Heine, W. 1943. Beobachtungen und experimentelle Untersuchungen über Manganvergiftungen und "Manganpneumonien". Z. Hyg. Infektionskrank. 125: 3-76. (Ger.)

Hem, J.O. 1970. Study and interpretation of the chemical characteristics of natural waters. U.S. Geol. Sur. Paper 1473, Washington, DC. p. 126-131.

Hewett, D.F. 1932. Manganese in sediments. In: Treatise on Sedimentation, W.H. Twenhofel, Ed. Williams and Wilkins Co., Baltimore, MD. p. 562-581.

Hidy, G.M., G. Lauer and S. Heisler. 1977. Assessment of the catalytic activity of manganese for SO<sub>2</sub> oxidation in the ambient air. Document P-4229. Prepared for Ethyl Corporation by Environmental Research and Technology, Inc., Westlake Village, CA.

Hill, R.M. and D.E. Holtkamp. 1954. Storage of dietary manganese and thiamine in the rat. J. Nutr. 53: 73-82.

Hillebrand, B.L. and K.L. Beattie. 1979. New method for the detection of misincorporation by DNA polymerases. Genetics. 91: S49.

Hirschler, D.A. and L.F. Gilbert. 1964. Nature of lead in automobile exhaust gas. Arch. Environ. Health. 8: 297.

Hodgson, J.F. 1963. Chemistry of the micronutrient elements. Advan. Agron. 15: 119-159.

Holbrook, D.J., Jr., M.E. Washington, H.B. Leake and P.E. Brubaker. 1975. Studies on the evaluation of the toxicity of various salts of lead, manganese, platinum and palladium. Environ. Health. Perspect. 10: 95-101.

Holtkamp, D.E. and R.M. Hill. 1950. The effect on growth of the level of manganese in the diet of rats, with some observations on the manganese-thiamine relationship. J. Nutr. 41: 307-316.

Hopps, H.C. 1977. The biologic bases for using hair and nail for analyses of trace elements. Sci. Total Environ. 7: 71-89.

Horiguchi, S., T. Utsonomiya, T. Kasahara, et al. 1966. A survey of the actual conditions of factories handling manganese compounds. Jap. J. Ind. Health. 8: 333-342. (Jap.)

Horiuchi, K., S. Horiguchi, N. Tanaka and K. Shinagawa. 1967. Manganese contents in the whole blood, urine and feces of a healthy Japanese population. Osaka City Med. J. 13: 151-163.

Horiuchi, K., S. Horiguchi, K. Shinagawa, T. Utsunomiya and Y. Tsuyama. 1970. On the significance of manganese contents in the whole blood and urine of manganese handlers. Osaka City Med. J. 16: 29-37.

Hoshishima, K., H. Tujii and K. Kano. 1978. Effects of the administration of trace amounts of metals to pregnant mice upon the behavior and learning of their offspring. Proc. Int. Congr. Toxicol. 1st. (1977), p. 569-570.

Hrsak, J. and M. Fugas. 1980. Preparation of samples of airborne particles for analysis of metals. Paper submitted for presentation at the Int. Microchemical Symp., Graz.

Hughes, E.R., S. Miller and G. Cotzias. 1966. Tissue concentrations of manganese and adrenal function. Am. J. Physiol. 211: 207-210.



Hurley, L.S. 1968. Approaches to the study of nutrition in mammalian development. Fed. Proc. 27: 193.

Hurley, L.S. 1981. The roles of trace elements in fetal and neonatal development. Philos. Trans. R. Soc. Lond. (Biol.) 294: 145-152.

Hurley, L.S. and L. Theriault-Bell. 1974. Genetic influence on response to dietary manganese deficiency in mice. J. Nutr. 104: 133-137.

Hutchinson, G.E. 1957. A Treatise on Limnology, Vol. 1. Geography, Physics, and Chemistry. John Wiley and Sons, Inc., New York. p. 1015.

IARC (International Agency for Research on Cancer). 1982. Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Vol. 29. Lyon, France.

ILO (International Labour Office). 1980. Occupational exposure limits for airborne toxic substances. A tabular compilation of values from selected countries. Occupational Safety and Health Series No. 37, Second ed., ILO, Geneva.

Imam, Z. and S.V. Chandra. 1975. Histochemical alterations in rabbit testis produced by manganese chloride. Toxicol. Appl. Pharmacol. 32: 534-544.

Instrumentation for Environmental Monitoring. 1975. Instrumentation for Environmental Monitoring, Air Part 2, First ed. Lawrence Berkley Lab., Univ. of California.

Issels, R.D. and E. Lengfelder. 1981. Superoxide dismutase activity in human lymphatic cells and in malignant lymphoma. *Exp. Hematol.* 9(S9): 184. (Abstr.)

Iversen, S.D. and L.L. Iversen. 1975. *Behavioral Pharmacology*. Oxford University Press, New York and Oxford.

Jacobs, R.M., A.O.L. Jones, B.E. Fry, Jr. and M.R. Spivey-Fox. 1970. Decreased long-term retention of  $^{115m}\text{Cd}$  in Japanese quail produced by a combined supplement of zinc, copper, and manganese. *J. Nutr.* 108: 901-910.

Järvinen, R. and A. Ahlström. 1975. Effect of the dietary manganese level on tissue manganese, iron, copper and zinc concentrations in female rats and their fetuses. *Med. Biol.* 53: 93-99.

Jindrichova, J. 1969. Anwendungsmöglichkeit der manganbestimmung im Stuhl ab Expositionstest. *Int. Arch. Gewerbepath. Gewerbehyg.* 25: 347-359. (Ger.)

Jonderko, G. 1965. Calcium, magnesium, inorganic phosphorus, sodium, potassium and iron levels in blood serum in the course of acute experimental manganese poisoning. *Med. Pr.* 16: 288-292.

Jonderko, G., A. Kujawska and H. Langaher-Lewowicka. 1971. Studies on the early symptoms of manganese toxicity. *Med. Pr.* 22: 1-10. (Pol.)

Jonderko, G., D. Dzekanska, T. Twardowski and E. Tyrna. 1973. Effects of occupational exposure to manganese on the development of atherosclerosis. Med. Pr. 24: 589-599. (Engl. Abstr.)

Jonderko, G., A. Kujawska and H. Langaher-Lewowicka. 1974. Effect of interruption of occupational contact with manganese upon neurological and biochemical symptoms of the toxic effects of manganese. Med. Pr. 25: 543-548.

Jones, J.M., J.T. McCall and L.R. Elvbeck. 1972. Trace metals in human osteogenic sarcoma. Mayo Clin. Proc. 47: 476-478.

Jones, T.S. 1982. Manganese. In: Minerals Yearbook 1981. Vol. 1. Metal and Minerals. U.S. Bureau of Mines, U.S. Dept. of Interior, Washington, DC.

Jorgenson, T.Z., C.J. Rushbrook, G.W. Hewell and S. Green. 1978. In vivo mutagenesis investigations of four GRAS chemicals. Mutat. Res. 53: 205. (Abstr.)

Josephson, J. 1980. Safeguards for groundwater. Environ. Sci. Technol. 14: 38-44.

Jötten, K.W., H. Reploh and G. Hegemann. 1939. Experimentelle Untersuchungen über die Manganpneumonie und ihre Beziehungen zur Thomas-schlackenpneumonie. Arch. Gewerbepath. Gewerbehyg. 9: 314-336. (Ger.)

Juday, C., E.A. Birge and V.W. Meloche. 1933. Mineral content of the lake waters of northeastern Wisconsin. Wisconsin Acad. Sci. Arts Lett. Trans. 31: 223-275.

Jungers, R.J. 1973. Manganese in motor oil. U.S. EPA, Natl. Res. Center, Research Triangle Park, NC, May 1973. (Unpubl.)

Kagamimori, S., I. Makino, Y. Hiramaru, et al. 1973. Epidemiological studies on disturbance of respiratory system caused by manganese air pollution (report 2). Improvement of respiratory functions after setting up dust collectors. Jap. J. Pub. Health. 20: 413-421.

Kato, M. 1963. Distribution and excretion of radiomanganese administered to the mouse. Q. J. Exp. Physiol. 48: 355-369.

Kaur, G., S.K. Hasan and R.C. Srivastava. 1980. The distribution of manganese-54 in fetal, young and adult rats. Toxicol. Lett. 5: 423-426.

Kehoe, R.A., J. Cholak and R.V. Story. 1940. A spectrochemical study of the normal ranges of concentration of certain trace metals in biological materials. J. Nutr. 19: 579-592.

Kemmerer, A.R., C.A. Elvehjem and E.B. Hart. 1931. Studies on the relation of manganese to the nutrition of the mouse. J. Biol. Chem. 92: 623-630.

Kesic, B. and V. Häusler. 1954. Hematological investigation on workers exposed to manganese dust. Arch. Ind. Hyg. Occup. Med. 10: 336-343.

Khavtasi, A.A. 1958. The possibility of chronic manganese poisoning among workers in manganese mines. Gig. Tr. Prof. Zabol. 2: 36-39.

Khazan, G.L., Y.M. Stanislavskiy, Y.V. Vasilenko, et al. 1956. Working conditions and health status of workers in the Nikopol's manganese mines. Vrach. Delo. 1-5: 277-291.

Kimura, M., N. Yagi and Y. Itokawa. 1978. Effect of subacute manganese feeding on serotonin metabolism in the rat. J. Toxicol. Environ. Health. 4: 701-707.

King, D.L. 1971. Sampling in natural waters and waste effluents. In: Water and Water Pollution Handbook, Vol. 2, L.L. Clacco, Ed. M. Dekker, Inc., New York. p. 451-481.

Klaassen, C.D. 1974. Biliary excretion of manganese in rats, rabbits, and dogs. Toxicol. Appl. Pharmacol. 29: 458-468.

Klawans, H., Jr., M.M. Haki and D. Shenker. 1970. Theoretical implications of the use of L-dopa in Parkinsonism. Acta Neurol. Scand. 46: 409-411.

Kleinkopf, M.D. 1960. Spectrographic determination of trace elements in lake waters of northern Maine. Geol. Soc. Am. Bull. 71: 1221-1242.

Kleinman, M.T., T.J. Kneip, D.M. Bernstein and M. Eisenbud. 1975. Fallout of toxic trace metals in New York City. Proc. 15th Annual Hanford Life Science Symposium on Biological Implications of Metals in the Environment, Seattle, WA.

Kleinman, M.T., G.S. Pasternack, M. Eisenbud and T.J. Kneip. 1980. Identifying and estimating the relative importance of sources of airborne particulates. Environ. Sci. Technol. 14: 62-65.

Klinkhammer, G.P. 1980. Determination of manganese in sea water by flameless atomic absorption spectrometry after preconcentration with 8-hydroxyquinoline in chloroform. Anal. Chem. 52: 117-120.

Kneip, T.J., R.P. Mallon and M.T. Kleinman. 1983. The impact of changing air quality on multiple regression models for coarse and fine particle fractions. Atmos. Environ. 17: 299-304.

Konetzka, W.A. 1977. Microbiology of metal transformations. In: Microorganisms and Minerals, E.O. Weinberg, Ed. Marcel Dekker, Inc., New York. p. 317-342.

Kontur, P. and L. Fechter. 1983. A neurochemical and behavioral study on neonatal manganese exposure. Neurotoxicology. 3: 143-144.

Kopp, J.F. and R.C. Kroner. 1969. Trace metals in waters of the United States. A five-year summary of trace metals in rivers and lakes of the United States (Oct. 1, 1962-Sept. 30, 1967). Div. Pollut. Surveillance, Fed. Water Pollut. Control Admin., U.S. Dept. Interior, Cincinnati, OH. 32 p.

Kosai, M.F. and A.J. Boyle. 1956. Ethylenediaminetetraacetic acid in manganese poisoning of rats. A preliminary study. Ind. Med. Surg. 25: 1-3.

Koshida, Y., M. Kato and T. Hara. 1963. Autoradiographic observations of manganese in adult and embryo mice. *Q. J. Exp. Physiol.* 48: 370-378.

Kostial, K., D. Kello, S. Jugo, I. Rabar and T. Maljkovic. 1978. Influence of age on metal metabolism and toxicity. *Environ. Health Perspect.* 25: 81-86.

Kostial, K., I. Rabar, M. Blanusa and I. Simonovic. 1980. The effect of iron additive to milk on cadmium, mercury and manganese absorption in rats. *Environ. Res.* 22: 40-45.

Kuznetsov, S.I. 1970. *The Microflora of Lakes.* University of Texas Press, Austin. p. 365-407.

Lai, J.C.K. 1983. Personal communications.

Lai, J.C.K., T.K.C. Leung, J.F. Guest, L. Lim and A.N. Davison. 1980. The monoamine oxidase inhibitors clorgyline and 1-deprenyl also affect the uptake of dopamine, noradrenaline and serotonin by rat brain synaptosomal preparations. *Biochem. Pharmacol.* 29: 2763-2767.

Lai, J.C.K., T.K.C. Leung and L. Lim. 1981a. Brain regional distribution of glutamic acid decarboxylase, choline acetyltransferase, and acetylcholinesterase in the rat: Effects of chronic manganese chloride administration after two years. *J. Neurochem.* 36: 1443-1448.

La1, J.C.K., M.J. Minski, A.W.K. Chan, L. Lim and A.N. Davison. 1981b. Brain regional manganese distribution after chronic manganese treatment. *Biochem. Soc. Trans.* 9: 228.

La1, J.C.K., L. Lim and A.N. Davison. 1981c. Differences in the inhibitory effect of  $\text{Cd}^{2+}$ ,  $\text{Mn}^{2+}$  and  $\text{Al}^{3+}$  on the uptake of dopamine by synaptosomes from forebrain and from striatum of the rat. *Biochem. Pharmacol.* 30: 3123-3125.

La1, J.C.K., T.K.C. Leung, J.F. Guest, A.N. Davison and L. Lim. 1982a. The effects of chronic manganese chloride treatment expressed as age-dependent, transient changes in rat brain synaptosomal uptake of amines. *J. Neurochem.* 38: 844-847.

La1, J.C.K., T.K.C. Leung and L. Lim. 1982b. Monoamine oxidase activities in liver, heart, spleen and kidney of the rat. Organ-specific changes in aging and after chronic manganese chloride administration. *Exp. Geront.* 17: 219-225.

La1, J.C.K., T.K.C. Leung and L. Lim. 1982c. The ontogeny of acetylcholinesterase activities in rat brain regions and the effect of chronic treatment with manganese chloride. *J. Neurochem.* 39: 1767-1769.

La1, J.C.K., P.C.L. Wong and L. Lim. 1983a. Structure and function of synaptosomal and mitochondrial membranes: Elucidation using neurotoxic metals and neuromodulatory agents. *In: Neural Membranes*, G.Y. Sun, A.Y. Sun and J.-Y. Wu, Ed. Humana Press, Inc., Clifton, NJ.



Lai, J.C.K., T.K.C. Leung and L. Lim. 1983b. Differences in the neurotoxic effects of manganese during development and aging: Some observations on brain regional neurotransmitter and non-neurotransmitter metabolism in a developmental rat model of chronic manganese encephalopathy. *Neurotoxicology* (Mn Special Issue) (In press)

Lai, J.C.K., T.K.C. Leung, L. Lim, A.W.K. Chan and M.J. Minski. 1983c. Effects of chronic manganese treatment on rat brain regional sodium-potassium-activated and magnesium-activated adenosine triphosphatase activities during brain development. *Neurotoxicology* (Mn Special Issue) (In press)

Lai, J.C.K., A. Baker and J.P. Blass. 1983d. Differential inhibitory effects of metal ions on brain hexokinase. *Fed. Proc.* 42: 627.

Lai, J.C.K., A.W.K. Chan, M.J. Minski, T.K.C. Leung, L. Lim and A.N. Davison. 1983e. In: Metal Ions in Neurology and Psychiatry, S. Gabay, J. Harris and B.T. Ho, Ed. Alan R. Liss, Inc., New York. (in preparation)

Langmyhr, F.J., T. Lind and J. Jonsen. 1975. Atomic absorption spectrometric determination of manganese, silver and zinc in dental material by atomization directly from the solid state. *Anal. Chim. Acta.* 80: 297-301.

Larsen, N.A., H. Pakkenberg, E. Damsgaard and K. Heydorn. 1979. Topographical distribution of arsenic, manganese, and selenium in the normal human brain. *J. Neurol. Sci.* 42: 407-416.

Laskey, J.W., G.L. Rehnberg, J.F. Hein and S.D. Carter. 1982. Effects of inorganic manganese ( $Mn_3O_4$ ) exposure on selected reproductive parameters in rats. J. Toxicol. Environ. Health. 9: 677-687.

Lassiter, J.W., W.J. Miller, M.W. Neathery, et al. 1974. Manganese metabolism and homeostasis in calves and rats. In: Trace Element Metabolism in Animals-2, W.G. Hoekstra, J.W. Suttie, M.E. Ganther and W. Mertz, Ed. University Park Press, Baltimore, Maryland. p. 557-559.

Lazrus, A.L., E. Lorange and J.P. Lodge, Jr. 1970. Lead and other metal ions in U.S. precipitation. Environ. Sci. Technol. 4: 55-58.

Leach, R.M., Jr. 1971. Role of manganese in mucopolysaccharide metabolism. Fed. Proc. 30: 991-994.

Leach, R.M., Jr. 1976. Metabolism and function of manganese. In: Trace Elements in Human Health and Disease, A.S. Prasad and D. Oberleas, Ed. Academic Press, New York. 2: 235-247.

Leach, R.M., Jr. and M.S. Lilburn. 1978. Manganese metabolism and its function. World Rev. Nutr. Diet. 32: 123-134.

Lee, C., N.B. Kim, I.C. Lee and K.S. Chung. 1977. The use of a chelating resin column for preconcentration of trace elements from sea water in their determination by neutron activation analysis. Talanta. 24: 241-245.

Lee, R.E., S.S. Goranson, R.E. Enrione and G.B. Morgan. 1972. The National Air Surveillance Cascade Impactor Network. Part II. Size distribution of trace-metal components. Environ. Sci. Technol. 6: 1025-1030.

Lentsch, J.W., T.J. Kneip, M.E. Wrenn, G.P. Howells and M. Eisenbud. 1972. Stable manganese and  $Mn^{54}$  distribution in the physical and biological components of the Hudson River estuary. Proc. Third Natl. Symp. on Radioecol., Oak Ridge, TN.

Leung, T.K.C., J.C.K. Lai and L. Lim. 1981. The regional distribution of monoamine oxidase activities towards different substrates: Effects in rat brain of chronic administration of manganese chloride and of aging. J. Neurochem. 36: 2037-2043.

Leung, T.K.C., J.C.K. Lai and L. Lim. 1982a. The effects of chronic manganese feeding on the activity of monoamine oxidase in various organs of the developing rats. Comp. Biochem. Physiol. 71C: 223-228.

Leung, T.K.C., J.C.K. Lai, M. Tricklebank, A.N. Davison and L. Lim. 1982b. Chronic manganese treatment of rats alters synaptosomal transport of dopamine and the behavioral response to amphetamine administration. J. Neurochem. 39: 1496-1499.

Levina, Z.N. and E.G. Robacovskaja. 1955. Changes in the lung tissues under intratracheal injections of manganese oxides. Gig. Sanit. 20: 25-28.

Lewis, C.W. and E.S. Macias. 1980. Composition of size-fractionated aerosol in Charleston, West Virginia. Atmos. Environ. 14: 185-194.

Linder, H.R., H.D. Seltner and B. Schreiber. 1978. Use of dibenzylidithiocarbamate as coprecipitant in the routine determination of 12 heavy metals in pharmaceutical by X-ray fluorescence spectroscopy. Anal. Chem. 50: 896-899.

Liu, C.-K., B.A. Roscoe, K.G. Severin and P.K. Hopke. 1982. The application of factor analysis to source apportionment of aerosol mass. Am. Ind. Hyg. Assoc. J. 43: 314-318.

Lloyd-Davies, T.A. 1946. Manganese pneumonitis. Br. J. Ind. Med. 3: 111-135.

Lloyd-Davies, T.A. and H.E. Harding. 1949. Manganese pneumonitis. Further clinical and experimental observations. Br. J. Ind. Med. 6: 82-90.

Loven, D.P., L.W. Oberly, F.M. Rousseau and K.H. Stevens. 1980. Superoxide dismutase activity in 1,2-dimethylhydrazine-induced rat colon adenocarcinoma. J. Natl. Cancer Inst. 65: 377-381.

Luthy, R.G. 1964. New concept for iron bacteria control in water wells. Water Well J. 24: 24, 29-30.

Mahomedy, M.C., Y.H. Mahomedy, P.A.S. Canhan, J.W. Downing and D.E. Jeal. 1975. Methaemoglobinaemia following treatment dispensed by witch doctors. Anaesthesia. 30: 190-193.

Mahoney, J.P. and W.J. Small. 1968. Studies on manganese. III. The biological half-life of radiomanganese in man and factors which affect this half-life. J. Clin. Invest. 47: 643-653.

Mahoney, J.P., K. Sargent, M. Greland and W.J. Small. 1969. Studies on manganese. I. Determination in serum by atomic absorption spectrophotometry. Clin. Chem. 15: 312-322.

Maigetter, R.Z., R. Ehrlich, J.D. Fenters and C.E. Gardner. 1976. Potentiating effect of manganese dioxide on experimental respiratory infections. Environ. Res. 11: 386-391.

Marjanen, H. 1969. Possible causal relationship between the easily soluble amount of manganese on arable mineral soil and susceptibility to cancer in Finland. Ann. Agric. Fenn. 8: 326-334.

Massaro, E.J., R.B. D'Agostino, C. Stineman, J.B. Morganti and B.A. Lown. 1980. Alterations in behavior of adult offspring of female mice exposed to  $MnO_2$  dust during gestation. Fed. Proc. Fed. Am. Soc. Exp. Biol. 39: 623. (Abstr.)

Matousek, J. and B. Stevens. 1971. Biological applications of the carbon-rod atomizer in A.A.S. preliminary studies on Mg, Fe, Cu, Pb, Zn in blood and plasma. Clin. Chem. 17: 363.

Matricardi, L.R. and J.H. Downing. 1981. Manganese and manganese alloys. In: Kirk-Othmer Encyclopedia of Chemical Technology, Third ed., Vol. 14, M. Grayson and D. Eckroth, Ed. John Wiley and Sons, Inc., New York. p. 824-843.

Matrone, G., R.H. Hartman and A.J. Clawson. 1959. Studies of a manganese iron antagonism in the nutrition of rabbits and baby pigs. J. Nutr. 67: 309-317.

Matteson, M.J., W. Stober and H. Luther. 1969. Kinetics of the oxidation of sulfur dioxide by aerosols of manganese sulfate. Ind. Eng. Chem. Fundam. 8: 677-687.

Maynard, L.S. and G.C. Cotzias. 1955. The partition of manganese among organs and intracellular organelles of the rat. J. Biol. Chem. 214: 489-495.

Maynard, L.S. and S. Fink. 1956. The influence of chelation on radiomanganese excretion in man and mouse. J. Clin. Invest. 35: 831-836.

McCarty, M.F. 1981. Optimized mitochondrial function as a nutritional strategy in cancer immunotherapy. Med. Hypotheses. 7: 55-60.

McLeod, B.E. and M.F. Robinson. 1972. Dietary intake of manganese by New Zealand infants during the first six months of life. Br. J. Nutr. 27: 229-232.

McMullen, T.B., R.B. Faoro and G.B. Morgan. 1970. Profile of pollutant fractions in nonurban suspended particulate matter. J. Air Pollut. Control Assoc. 20: 369-372.

Hella, H. 1924. The experimental production of basal ganglion symptomatology in macacus rhesus. Arch. Neurol. Psychiatr. 11: 405-417.

Mena, I., O. Marin, S. Fuenzalida and G.C. Cotzias. 1967. Chronic manganese poisoning: Clinical picture and manganese turnover. Neurology. 17: 128-136.

Mena, I., K. Horiuchi, K. Burke and G.C. Cotzias. 1969. Chronic manganese poisoning. Individual susceptibility and absorption of iron. Neurology. 19: 1000-1006.

Mena, I., J. Court, S. Fuenzalida, P.S. Papavasiliou and G.C. Cotzias. 1970. Modification of chronic manganese poisoning. Treatment with L-dopa or 5-OH tryptophane. New Engl. J. Med. 282: 5-10.

Mena, I., K. Horiuchi and G. Lopez. 1974. Factors enhancing entrance of manganese into the brain: Iron deficiency and age. J. Nucl. Med. 15: 516.

Miller, S.T. and G.C. Cotzias. 1977. Effects on maternal tissue and milk manganese on the suckling mouse. Gen. Physiol.

Miller, S.T., G.C. Cotzias and H.A. Evert. 1975. Control of tissue manganese: Initial absence and sudden emergence of excretion in the neonatal mouse. Am. J. Physiol. 229: 1080-1084.

Miller, W.J., M.W. Neather, R.P. Gentry, D.M. Blackmon, J.W. Lassiter and F.M. Pate. 1972. Distribution and turnover rates of radioactive manganese in various tissues after duodenal dosing in Holstein calves fed a practical-type diet. J. Anim. Sci. 34: 460-464.

Mitchell, R.L. 1971. Trace elements in soils. G.T. Br. Min. Agric. Fish Food, Tech. Bull. No. 21. p. 8-20.

Mokievskaya, V.V. 1961. Manganese in water of the Black Sea. Dokl. Nauk SSSR. 137: 1445-1447.

Moody, J.R. and R.M. Lindstrom. 1977. Section and cleaning of plastic containers for storage of trace element samples. Anal. Chem. 49: 2264-2267.

Moore, F.L. 1983a. Letter from F.L. Moore, Elkem Metals Co., Marietta, OH to L.S. Erdreich, Environmental Criteria and Assessment Office, U.S. EPA, Cincinnati, OH. February 22.

Moore, F.L. 1983b. Letter from F.L. Moore, Elkem Metals Co., Marietta, OH to R.J.F. Bruins, Environmental Criteria and Assessment Office, U.S. EPA, Cincinnati, OH. March 8.

Moore, W., J.F. Stara, D. Hysell, M. Malanchuk, J. Burkart and R. Hinners. 1975. Toxicologic evaluations of fuel additive -- Methylcyclopentadienyl Manganese Tricarbonyl (MMT). Presented at the 1975 SAE Automobile Engineering Meeting, Detroit, MI, October 13-17.



Morel, F. and J.J. Morgan. 1972. Numerical method for computing equilibriums in aqueous chemical system. Environ. Sci. Technol. 6: 58-67.

Morgan, J.J. 1967. Chemical equilibriums and kinetic properties of manganese in natural waters. In: Principles and Applications of Water Chemistry, S.D. Faust and J.V. Hunter, Ed. John Wiley and Sons, Inc., New York. p. 561-624.

Morgan, J.M. 1972. Hepatic copper, manganese and chromium content in bronchogenic carcinoma. Cancer. 29: 710-713.

Mouri, T. 1973. Experimental study on the inhalation of manganese dust. Shikoku Acta Med. 28: 118-129.

Mueller, P.K., W.K. Gong and E.E. Jones. 1963. Aerosol sampling methods applied to auto exhaust. Fifth Conf. "Methods in Air Pollution Studies", California State Dept. Pub. Health, Los Angeles, CA.

Mulay, I.L., R. Roy, B.E. Knox, N.H. Suhr and W.E. Delaney. 1971. Trace metal analysis of cancerous and non-cancerous human tissues. J. Natl. Cancer Inst. 47: 1-13.

Murthy, G.K., U.S. Rhea and J.T. Peeler. 1971. Levels of antimony, cadmium, chromium, cobalt, manganese and zinc in institutional total diets. Environ. Sci. Technol. 5: 436-442.

Murthy, R.C., S. Lal, D.K. Saxena, G.S. Shukla, M.M. Ali and S.V. Chandra. 1981. Effect of manganese and copper interaction on behavior and biogenic amines in rats fed a 10% casein diet. *Chem. Biol. Interact.* 37: 299-308.

Musani-Marazovic, Lj. and Z. Pucar. 1977. Interaction of  $^{54}\text{Mn}$  and  $^{55(59)}\text{Fe}$  with EDTA in seawater and 0.55 M NaCl solution. *Mar. Chem.* 5: 229-242.

Mustafa, S.J. and S.V. Chandra. 1971. Levels of 5-hydroxytryptamine, dopamine and norepinephrine in whole brain of rabbits in chronic manganese toxicity. *J. Neurochem.* 18: 931-933.

Mustafa, S.J. and S.V. Chandra. 1972. Adenosine deaminase and protein pattern in serum and cerebrospinal fluid in experimental manganese encephalopathy. *Arch. Toxicol.* 28: 279-285.

Muzzarelli, R.A.A. and R. Rochetti. 1975. Atomic-absorption determination of manganese, cobalt and copper in whole blood and serum, with a graphite atomizer. *Talanta.* 22: 683-685.

NAPCA (National Air Pollution Control Administration). 1970. Kanawha Valley Air Pollution Study. Research Triangle Park, NC. Publ. No. APTD 70-1, March. p. 367.

NAS (National Academy of Sciences). 1973. Medical and Biologic Effects of Environmental Pollutants: Manganese. NAS, Washington, DC. p. 126-131.

NAS (National Academy of Sciences). 1978. Nutrient requirements of laboratory animals, Third ed. NAS, Washington, DC. 96 p.

NASN (National Air Surveillance Network). 1957-1969. National Air Surveillance Network: Manganese data. National Aerometric Data Bank. U.S. EPA, Off. Air Qual. Plan. Stand., Research Triangle Park, NC.

Nazif, M. 1936. Manganese as an industrial poison. J. Egypt. Pub. Health Assoc. 10: 1-20.

NCI (National Cancer Institute). 1982. Notice of Research Project Tox-Tips. p. 72-79.

Neff, N.H., R.E. Barrett and E. Costa. 1969. Selective depletion of caudate nucleus dopamine and serotonin during chronic manganese dioxide administration to squirrel monkeys. Experientia. 25: 1140-1141.

Newberne, P.M. 1973. Input and disposition of manganese in man. In: Medical and Biologic Effects of Environmental Pollutants: Manganese. NAS, Washington, DC. p. 77-82.

Nichol, I., R.F. Horsnail and J.S. Webb. 1967. Geochemical patterns in stream sediment related to precipitation of manganese oxides. Trans. Inst. Min. Metall. Sect. B. 76: 113-115.

Nishiyama, K., Y. Suzuki, N. Fujii, H. Yano, T. Miyai and K. Ohmishi. 1975. Effect of long-term inhalation of manganese dusts. II. Continuous observation of the respiratory organs of monkeys and mice. Jap. J. Hyg. 30(1): 117.

Nogawa, K., E. Kobayashi, M. Sakamoto, et al. 1973. Epidemiological studies on disturbance of respiratory system caused by manganese air pollution. Report 1: Effects on respiratory system of junior high school students. Jap. J. Pub. Health. 20: 315-326. (Japanese with English abstract)

North, B.B., J.M. Leichsenring and L.M. Norris. 1960. Manganese metabolism in college women. J. Nutr. 72: 217-223.

Oberleas, D. and D.F. Caldwell. 1981. Trace minerals in pregnancy. Int. J. Environ. Stud. 17: 85-98.

Oberly, L.W. and G.R. Buettner. 1979. Role of superoxide dismutase in cancer: A review. Cancer Res. 39: 1141-1149.

Oberly, L.W., S.K. Saku and F.J. Cheng. 1978. Superoxide dismutase activity in normal mouse brain, prenatal brain and neuroblastoma cells. Proc. Am. Assoc. Cancer Res. 19: 147. (Abstr.)

Oberly, L.W., T.D. Oberly and G.R. Buettner. 1980. Cell differentiation, aging and cancer: The possible roles of superoxide; and superoxide dismutases. Med. Hypotheses. 6: 240-268.

Olahy, D.A., R.A. Schmitt and W.F. Bethard. 1966. Neutron activation analysis of magnesium, calcium, strontium, barium, manganese, cobalt, copper, zinc, sodium and potassium in human erythrocytes and plasma. J. Nucl. Med. 7: 917-927.

Ondov, J.M., R.C. Ragaini and A.H. Biermann. 1979. Emissions and particle-size distributions of minor and trace elements at two Western coal-fired power plants equipped with cold-side electrostatic precipitators. Environ. Sci. Technol. 13: 946-953.

Onoda, K., A. Hasegawa, M. Sunouchi, et al. 1978. Studies on the fate of poisonous metals in experimental animal (VII). Distribution and transplacental passage of manganese in pregnant rat and fetus. J. Food Hyg. Soc. 19(2): 208-215.

Orent, E.R. and E.V. McCollum. 1931. Effects of deprivation of manganese in the rat. J. Biol. Chem. 92: 651-670.

Pace, T.G. and M.H. Frank. 1983. Procedures for estimating probability of non-attainment of a  $PM_{10}$  NAAQS using total suspended particulate or inhalable particulate data. Office of Air Quality Planning and Standards, Monitoring and Data Analysis Division, U.S. EPA, Research Triangle Park, NC. March.

Papavasiliou, P.S. and G.C. Cotzias. 1961. Neutron activation analysis: The determination of manganese. J. Biol. Chem. 236: 2365-369.

Papavasiliou, P.S., S.T. Miller and G.C. Cotzias. 1966. Role of liver in regulating distribution and excretion of manganese. *Am. J. Physiol.* 211(1): 211-216.

Parkinson, T.F., L.E. Millikan and D.F. Anderson. 1979. Manganese, copper and zinc concentrations in human skin lesions. *J. Appl. Radiation Isotop.* 30: 411-415.

Parnitzke, K.H. and J. Pfeiffer. 1954. On the clinical and pathological anatomy in chronic manganese poisoning. *Arch. Psychiatr. Z. Neurol.* 192: 405-429.

Paterni, L. 1954. Blood diseases from metals. *Folia Med.* 37: 994-1006.

Penalver, R. 1955. Manganese poisoning. The 1954 Ramazzini oration. *Ind. Med. Surg.* 24: 1-7.

Pentschew, A., F.F. Ebner and R.M. Kovatch. 1963. Experimental manganese encephalopathy in monkeys. *J. Neuropathol. Exp. Neurol.* 22: 488-499.

Perry, H.M., Jr., E.F. Perry, J.E. Purifoy and J.N. Erlanger. 1973. A comparison of intra- and interhepatic variability of trace metal concentrations in normal men. In: Trace Substances in Environmental Health. Proceedings University of Missouri 7th Annual Conference. University of Missouri, Columbia, MO. p. 281-288.

Pierson, W.R., D.E. McKee, W.W. Grachaczek and J.W. Butler. 1978. Methylcyclopentadienyl manganese tricarbonyl: Effect on manganese emissions from vehicles on the road. J. Air Pollut. Control Assoc. 28: 692-693.

Pilat, M.J., D.S. Ensor and J.C. Bosch. 1970. Source test cascade impactor. Atmos. Environ. 4: 671-679.

Piscator, M. 1976. Health hazards from inhalation of metal fumes. Environ. Res. 11: 268-270.

Piscator, M. and V. Vouk. 1979. Sampling and analytical methods. In: Handbook on the Toxicology of Metals, L. Friberg et al., Ed. Elsevier/North Holland Biomedical Press. p. 33-45.

Piver, W.T. 1974. Potential dilemma: The methods of meeting automotive exhaust emission standards of the Clean Air Act of 1970. Environ. Health Perspect. 8: 165-190.

Plumlee, M.P., D.M. Thrasher, W.M. Beesen, F.N. Andrews and H.E. Parker. 1956. The effects of a manganese deficiency upon the growth, development and reproduction of swine. J. Anim. Sci. 15: 352-367.

Pollack, S., J.N. George, R.C. Reba, R.M. Kaufman and W.H. Crosby. 1965. The absorption of nonferrous metals in iron deficiency. J. Clin. Invest. 44(9): 1470-1473.

Prasad, K.N., Ed. 1978. Trace Elements and Iron in Human Metabolism: Manganese. John Wiley and Sons, Inc., New York. p. 191-201.

Price, H.O. and G.E. Bunce. 1972. Effect of nitrogen and calcium on balance of copper, manganese and zinc in preadolescent girls. Nutr. Rep. Intl. 5: 275-280.

Rabar, I. 1976. Some factors influencing manganese metabolism in rats. M.Sc. Thesis, Univ. Zagreb, Zagreb, Yugoslavia.

Rall, D.P. 1969. Difficulties in extrapolating the results of toxicity studies in laboratory animals to man. Environ. Res. 2: 360-367.

Rehnberg, G.L., J.F. Hein, S.D. Carter and J.W. Laskey. 1980. Chronic manganese oxide administration to preweanling rats: Manganese accumulation and distribution. J. Toxicol. Environ. Health. 6: 217-226.

Rehnberg, G.L., J.F. Hein, S.D. Carter, R.S. Linko and J.W. Laskey. 1981. Chronic ingestion of  $Mn_3O_4$  by young rats: Tissue accumulation, distribution, and depletion. J. Toxicol. Environ. Health. 7: 263-272.

Rehnberg, G.L., J.F. Hein, S.D. Carter, R.S. Linko and J.W. Laskey. 1982. Chronic ingestion of  $Mn_3O_4$  by rats: Tissue accumulation and distribution of manganese in two generations. J. Toxicol. Environ. Health. 9: 175-188.

Reidies, A.H. 1981. Manganese Compounds. In: Kirk-Othmer Encyclopedia of Chemical Technology, Third ed., Vol. 14, M. Grayson and D. Eckroth, Ed. John Wiley and Sons, Inc., New York. p. 844-895.



Riley, J.P. and D. Taylor. 1968. Chelating resins for the concentration of trace elements from sea water and their analytical use in conjunction with atomic absorption spectrophotometry. *Anal. Chim. Acta.* 40: 479-485.

Robertson, D.E. and R. Carpenter. 1974. Neutron activation techniques for the measurement of trace metals in environmental samples. U.S. AEC, Tech. Info. Center, Off. Info. Ser., NAS-Ns-3174.

Rodier, J. 1955. Manganese poisoning in Moroccan miners. *Br. J. Ind. Med.* 12: 21-35.

Rodier, J. and M. Rodier. 1949. Le manganisme dans les mines marocaines. *Bull. Inst. Hyg. Maroc.* 9: 3-98. (Fre.)

Roldan, V. 1956. U. Manganismo profesional. Observaciones clinicas. In: Tercer Congreso Americano de Medicina del Trabajo, Vol. I., Caracas 1955. Union Americana de Medicine del Trabajo, Buenos Aires. p. 100-104.

Rosenstock, H.A., D.G. Simons and J.S. Meyer. 1971. Chronic manganism. Neurologic and laboratory studies during treatment with levo-dopa. *J. Am. Med. Assoc.* 217: 1354-1358.

Roussel, B. and B. Renaud. 1977. Effect of chronic manganese intoxication on the sleep-wake cycle in the rat. *Neurosci. Lett.* 4: 55-60.

Ruch, R.R., H.J. Gluskoter and N.F. Shimp. 1973. Occurrence and distribution of potentially volatile trace elements in coal. Interim Report, Jan.-Dec. 1972. EPA Contract 68-02-0246 by Illinois State Geol. Sur. Environ. Geol. Notes 61. 43 p.

Sabadas, E. 1969. On neurohumoral regulation of the daily cycle of the manganese level of the blood. In: II Vsesojuznyh biohimiceskij S'ezd g. Taskent 1969, Tez. seke. soobc. Tashkent, Izdatel'stvo FAN Uzbekskoj SSR. 13: 159-160. (Rus.)

Sabnis, C.V., P.K. Yennawar, V.L. Pampattiwar and J.M. Deshpande. 1966. An environmental study of a ferro-manganese alloy concern. Indian J. Ind. Med. 11: 207-222.

Santoliquido, P.M., H.W. Scuthwick and J.H. Olwin. 1976. Trace metal levels in cancer of the breast. Surg. Gynecol. Obstet. 142: 65-70.

Saric, M. 1978. Biological effects of manganese. U.S. EPA, Research Triangle Park, NC. EPA 600/1-78-001. p. 152.

Saric, M. 1983. Personal communication.

Saric, M. and O. Hrustic. 1975. Exposure to airborne manganese and arterial blood pressure. Environ. Res. 10: 314-318.

Saric, M. and S. Lucic-Palaic. 1977. Possible synergism of exposure to airborne manganese and smoking habit in occurrence of respiratory symptoms. In: Inhaled Particles, IV, W.H. Walton, Ed. Pergamon Press, New York. p. 773-779.

Saric, M., A. Holetic and E. Ofner. 1975. Acute respiratory diseases in a manganese contaminated area. In: Proceedings of the International Conference on Heavy Metals in the Environment. Toronto, Canada. p. 389-398.

Saric, M., A. Markicevic and O. Hrustic. 1977. Occupational exposure to manganese. Br. J. Ind. Med. 34: 114-118.

Saxena, J. and P.H. Howard. 1977. Environmental transformation of alkylated and inorganic forms of certain metals. Adv. Appl. Microbiol. 21: 185-226.

Scander, A. and H.A. Sallam. 1936. A report on eleven cases of chronic manganese poisoning. J. Egypt. Med. Assoc. 19: 57-62.

Scheuhammer, A.M. and M.G. Cherian. 1981. The influence of manganese on the distribution of essential trace elements. I. Regional distribution of Mn, Na, K., Mg, Zn, Fe, and Cu in rat brain after chronic Mn exposure. Toxicol. Appl. Pharmacol. 61: 227-233.

Schlage, C. and B. Wortberg. 1972. Manganese in the diet of healthy pre-school and school children. Acta Paediat. Scand. 61: 648-652.

Schopper, W. 1930. Über Lungenentzündungen bei Brauneisenstein-Bergarbeitern. Arch. Hyg. Bakt. 104: 175-183. (Ger.)

Schroeder, H.A. 1970. Manganese -- Air quality monograph. Am. Petroleum Inst., No. 70-17, Washington, DC. 34 p.

Schroeder, H.A. and A.P. Nason. 1976. Interactions of trace metals in mouse and rat tissues: Zinc, chromium, copper, and manganese with 13 other elements. J. Nutr. 106: 198-203.

Schroeder, H.A., D.D. Balassa and I.H. Tipton. 1966. Essential trace elements in man: Manganese, a study on homeostasis. J. Chron. Dis. 19: 545-571.

Schroeder, H.A., M. Mitchener and A.P. Nason. 1974. Life-term effects of nickel in rats: Survival tumors, interactions with trace elements and tissue levels. J. Nutr. 104: 239-243.

Schuler, P., H. Oyanguren, V. Maturana, et al. 1957. Manganese poisoning. Environmental and medical study at a Chilean mine. Ind. Med. Surg. 26: 167-173.

Scrutton, M.C., P. Griminger and J.C. Wallace. 1972. Pyruvate carboxylase: Bound metal content of the vertebrate liver enzyme as a function of diet. J. Biol. Chem. 247: 3305-3313.

Seth, P.K., N. Nagar, R. Husain and S.V. Chandra. 1973. Effects of manganese on rabbit testes. *Environ. Physiol. Biochem.* 3: 263-267.

Seth, P.K., J.S. Hong, C.D. Kilts and S.C. Bordy. 1981. Alteration of cerebral neurotransmitter receptor function by exposure of rats to manganese. *Toxicol. Lett.* 9: 247-254.

Shacklette, H.T., J.C. Hamilton, J.G. Boerngen and J.M. Bowles. 1971. Elemental composition of surficial materials in the conterminous United States. U.S. Geol. Sur. Paper 574-D, Washington, DC. 71 p.

Shanker, R., R.K.S. Dogra, A.P. Sahu and S.H. Zaidi. 1976. Experimental manganese lymphadenopathy in guinea pigs. *Arch. Toxicol.* 36: 151-157.

Snigan, S.A. and B.R. Vitvitskaya. 1971. Experimental substantiation of permissible residual concentration of potassium permanganate in drinking water. *Gig. Sanit.* 36: 15-18.

Shimkin, M.B. and G.D. Stoner. 1975. Lung tumors in mice: Application to carcinogenesis bioassay. *Adv. Cancer Res.* 21: 1-58.

Shrader, R.E. and G.J. Everson. 1968. Pancreatic pathology in manganese-deficient guinea pigs. *J. Nutr.* 94: 269.

Shukla, G.S. and S.V. Chandra. 1976. Manganese induced morphological and biochemical changes in the brain of iron deficient rats. *Ind. Health.* 14: 87-92.

Shukla, G.S. and S.V. Chandra. 1977. Levels of sulfhydryls and sulfhydryl-containing enzymes in brain, liver and testis of manganese treated rats. Arch. Toxicol. 37: 319-325.

Shukla, G.S., S. Singh and S.V. Chandra. 1978. The interaction between manganese and ethanol in rats. Acta Pharmacol. Toxicol. 43: 354-362.

Sigan, S.A. and B.R. Vitvickaja. 1971. Experimental substantiation of permissible residual concentrations of potassium permanganate in drinking water. Gig. Sanit. 36: 15-18.

Silbergeld, E.K. 1982. Current status of neurotoxicology, basic and applied. Trends Neurosci. 5: 291-294.

Sillen, L.G. 1961. The physical chemistry of sea water. Oceanogr. Am. Assoc. Adv. Sci. p. 549-581.

Silver, S. and P. Jasper. 1977. Manganese transport in microorganisms. In: Microorganisms and Minerals, E.D. Weinberg, Ed. Marcel Dekker, Inc., New York. p. 105-150.

Singh, J., R. Hussain, S.K. Tandon, P.K. Seth and S.V. Chandra. 1974. Biochemical and histopathological alterations in early manganese toxicity in rats. Environ. Physiol. Biochem. 4: 16-23.

Singh, J., S.V. Chandra and S.K. Tandon. 1975. Chelation in metal intoxication. II. In vitro and in vivo effect of some compounds on brain, liver and testis of rats treated with manganese sulfate. Bull. Environ. Contam. Toxicol. 14: 497-504.

Singh, J., J.L. Kaw and S.H. Zaidi. 1977. Early biochemical response pulmonary tissue to manganese dioxide. Toxicology. 8: 177-184.

Singh, S., G.S. Shukla, R.S. Srivastava and S.V. Chandra. 1979. The interaction between ethanol and manganese in rat brain. Arch. Toxicol. 41: 307-316.

Sitaramayya, A., N. Nagar and S.V. Chandra. 1974. Effect of manganese on enzymes in rat brain. Acta Pharmacol. Toxicol. 35: 185-190.

Sittig, M. 1976. Toxic Metal, Pollution Control and Worker Protection. Noyes Data Corp., Park Ridge, NJ. p. 184-203.

Skreb, Y., J. Racic and N. Hors. 1980. Cytotoxicity of selected mutagens ( $Pb^{2+}$ ,  $Cd^{2+}$ ,  $Mn^{2+}$ ) to cultured mammalian cells. Mutat. Res. 74: 241-242.

Slavin, S., W.B. Barnett and H.L. Kahn. 1972. The determination of atomic absorption detection limits by direct measurement. Atomic Absorp. Newslett. 11: 37-41.

Smeijers-Verbeke, J., P. Lowenthal and D.L. Massart. 1976. Distribution of Mn in fly ash. Clin. Chim. Acta. 68: 343-347.

Smith, R.D. and K.K. Nielson. 1979. Concentration dependence of volatilized elements in fly ash. Environ. Sci. Technol. 13: 553-558.

Smyth, C.P. Carpenter, C.S. Weil, U.C. Pozzani, J.A. Striegel and J.S. Hycum. 1969. Range-finding toxicity data: List VII. J. Am. Ind. Hyg. Assoc. 30: 470-476.

Smyth, L.T., R.C. Ruhf, N.E. Whitman and T. Dugan. 1973. Clinical manganese and exposure to manganese in the production and processing of ferromanganese alloy. J. Occup. Med. 15: 101-109.

Spencer, D.W. and P.G. Brewer. 1971. Vertical advection diffusion and redox potentials as controls on the distribution of manganese and other trace metals dissolved in waters of the Black Sea. J. Geophys. Res. 76: 5877-5892.

Stadler, H. 1936. Histopathology of the brain resulting from manganese poisoning. Z. Ges. Neurol. Psychiat. 154: 62-76.

Standard Methods for the Examination of Water and Wastewater. 1971. Thirteenth ed. Am. Pub. Health Assoc. p. 213.



Stara, J.F., W. Moore, Jr., C. Hysell, et al. 1973. Toxicology of methylcyclopentadienyl manganese tricarbonyl (MMT) and related manganese compounds emitted from mobile and stationary sources. Proc. 4th Ann. Conf. Environ. Toxicol. AMRL-TR-73-125, Paper No. 20. p. 251-270.

Stevens, R.K., T.G. Dzuby, G. Russwurm and D. Rickel. 1978. Sampling and analysis of atmospheric sulfates and related species. Atmos. Environ. 12: 55-68.

Stoeppler, M., P. Valenta and H.W. Nurnberg. 1979. Application of independent methods and standard materials: An effective approach to reliable trace and ultratrace analysis of metals and metalloids in environmental and biological matrices. Fresenius Z. Anal. Chem. 297: 22-34.

Stoner, G.D., M.B. Shimkin, M.C. Troxell, T.L. Thompson and L.S. Terry. 1976. Test for carcinogenicity of metallic compounds by the pulmonary tumor response in strain A mice. Cancer Res. 36: 1744-1747.

Stumm, W. and H. Billinski. 1972. Trace metals in natural waters: Difficulties of interpretation arising from our ignorance on their speciation. In: Advances in Water Pollution Research, S.H. Jenkins, Ed. Pergamon Press, Oxford. p. 39-49.

Sullivan, R.J. 1969. Air pollution aspects of manganese and its compounds. U.S. DHEW, Publ. No. APTD 69-39, NAPCA, Raleigh, NC.

Sumino, K., K. Hayakawa, T. Shibata and S. Kitamura. 1975. Heavy metals in normal Japanese tissues. Arch. Environ. Health. 30: 487-494.

Sun, A.S., A. Cederbaum and L.R. Wasserman. 1980. Superoxide and hydrogen peroxide generating enzyme activities in normal rat liver and hepatoma HC-252. Proc. Am. Assoc. Cancer Res. 21: 26. (Abstr.)

Sunderman, F.W., Jr., T.J. Lau and L.J. Cralley. 1974. Inhibitory effect of manganese upon muscle tumorigenesis by nickel subsulfide. Cancer Res. 34: 92-95.

Sunderman, F.W., Jr., K.S. Kasprzak, T.J. Lau, et al. 1976. Effects of manganese on carcinogenicity and metabolism of nickel subsulfide. Cancer Res. 36: 1790-1800.

Sunderman, F.W., M.C. Reid, P.R. Allpass and S.B. Taubman. 1980. Manganese inhibition of sarcoma induction by benzo(a)pyrene in Fischer rats. Proc. Am. Assoc. Cancer Res. 21: 72. (Abstr.)

Suzuki, Y. 1970. Concerning environmental pollution by manganese. Ind. Med. 12: 529-533.

Suzuki, Y. 1974. Studies on excessive oral intake of manganese. II. Minimum dose for manganese accumulation in mouse organ. Shikoku Acta Med. 30: 32-45.

Suzuki, Y., K. Nishiyama, M. Doi, T. Hirose and H. Shibata. 1960. Studies on chronic manganese poisoning. *Tokushima J. Exp. Med.* 7: 124-132.

Suzuki, Y., K. Nishiyama, Y. Suzuki, et al. 1973a. The effects of chronic manganese exposure on ferromanganese workers (Part 1). *Shikoku Acta Med.* 29: 412-424. (Japanese with English Abstract)

Suzuki, Y., K. Nishiyama, Y. Suzuki, et al. 1973b. The effects of chronic manganese exposure on ferromanganese workers (Part 2). *Shikoku Acta Med.* 29: 433-438. (Japanese with English Abstract)

Suzuki, Y., T. Mouri, Y. Suzuki, K. Nishiyama, N. Fujii and H. Yano. 1975. Study of subacute toxicity of manganese dioxide in monkeys. *Tokushima J. Exp. Med.* 22: 5-10.

Suzuki, Y., N. Fujii, H. Yano, T. Ohkita, A. Ichikawa and K. Nishiyama. 1978. Effects of the inhalation of manganese dioxide dust on monkey lungs. *Tokushima J. Exp. Med.* 25: 119-125.

Swaine, D.J. 1955. The Trace-element Content of Soils. Commonwealth Bureau of Soil Science Technical Communication 48. Herald Printing Works, York, England. 157 p.

Swaine, D.J. and R.L. Mitchell. 1960. Trace-element distribution in soil profiles. *J. Soil Sci.* 11: 347-368.

Tanaka, S. and J. Lieben. 1969. Manganese poisoning and exposure in Pennsylvania. Arch. Environ. Health. 19: 674-684.

Tandon, S.K., S.V. Chandra, J. Singh, R. Hussain and P.K. Seth. 1975. Chelation in metal intoxication. I. In vivo effect of chelating agents on liver and testis of manganese administered rats. Environ. Res. 9: 18-25.

Ter Haar, G.L., D.L. Lenane, J.N. Hu and M. Brandt. 1972. Composition, size and control of automotive exhaust particulates. J. Air Pollut. Control Assoc. 22: 39-46.

Ter Haar, G.L., M.E. Griffing, M. Brandt, D.G. Oberding and M. Kapron. 1975. Methylcyclopentadienyl manganese tricarbonyl as an antiknock: Composition and fate of manganese exhaust products. J. Air Pollut. Control Assoc. 25: 858-860.

TGNA (Task Group on Metal Accumulation). 1973. Accumulation of toxic metals with special reference to their absorption, excretion and biological half-times. Environ. Physiol. Biochem. 3: 65-107.

Thompson, R.J. 1979. Collection and analysis of airborne metallic elements. In: Ultratrace Metal Analysis in Biological Sciences and Environment (Adv. Chem. Ser. 172), T.H. Risby, Ed. Washington, DC. Am. Chem. Soc. p. 54-72.

Thompson, R.J., G.B. Morgan and L.J. Purdue. 1970. Analysis of selected elements in atmospheric particulate matter by atomic absorption. Atomic Absorp. Newslett. 9: 53-57.

Thomson, A.B.R. and L.S. Valberg. 1972. Intestinal uptake of iron, cobalt, and manganese in iron-deficient rat. *Am. J. Physiol.* 223: 1327-1329.

Thomson, A.B.R., D. Olatunbosun, L.S. Valberg and J. Ludwig. 1971. Interrelation of intestinal transport system for manganese and iron. *J. Lab. Clin. Med.* 78: 642-655.

Tichy, M. and M. Cikrt. 1972. Manganese transfer into the bile in rats. *Arch. Toxicol.* 29: 51-58.

Tichy, M., J. Havrdova and M. Cikrt. 1971. Manganese determination in urine by method of atomic absorption spectrophotometry. *Pracov. Lek.* 23: 336-341.

Tichy, M., M. Cikrt and J. Havrdova. 1973. Manganese binding in rat bile. *Arch. Toxicol.* 30: 227-236.

Timaskina, N.S., V.M. Cheburanova and N.P. Timakin. 1981. Nickel, zinc, copper, iron and manganese content in the blood plasma and RBC of the patients with bronchogenic carcinoma. *Vopr. Onkol.* 27: 66.

Timonin, M.I. and G.R. Giles. 1952. Effect of different soil treatment on microbial activity and availability of manganese in manganese-deficient soil. *J. Soil Sci.* 3: 145-155.

Tipton, I.H. and M.J. Cook. 1963. Trace elements in human tissue. Part II. Adult subjects from the United States. *Health Physics.* 9: 103-145.

Tipton, I.H., P.L. Stewart and J. Dickson. 1969. Patterns of elemental excretion in long term balance studies. *Health Physics*. 16: 455-462.

Trendtel, F. 1936. Zur Frage des Manganismus. *Monatschr. Unfallh.* 43: 69-84. (Ger.)

Tsalev, D.L., F.J. Langmyhr and N. Gunderson. 1977. Direct atomic absorption spectrometric determination of manganese in whole blood of unexposed individuals and exposed workers in a Norwegian manganese alloy plant. *Bull. Environ. Contam. Toxicol.* 17: 660-666.

Turekian, K.K. and K.H. Wedepohl. 1961. Distribution of the elements in some major units of the earth's crust. *Geol. Soc. Am. Bull.* 72: 175-192.

Ulrich, C.E., W. Rinehart and W. Busey. 1979a. Evaluation of the chronic inhalation toxicity of a manganese oxide aerosol. I. Introduction, experimental design, and aerosol generation methods. *Am. Ind. Hyg. Assoc. J.* 40: 238-244.

Ulrich, C.E., W. Rinehart, W. Busey and M.A. Dorato. 1979b. Evaluation of the chronic inhalation toxicity of a manganese oxide aerosol. II. Clinical observations, hematology, clinical chemistry and histopathology. *Am. Ind. Hyg. Assoc. J.* 40: 322-329.

Ulrich, C.E., W. Rinehart and M. Brandt. 1979c. Evaluation of the chronic inhalation toxicity of a manganese oxide aerosol. III. Pulmonary function, electromyograms, limb tremor, and tissue manganese data. *Am. Ind. Hyg. Assoc. J.* 40: 349-353.

Underwood, E.J., Ed. 1977. Trace Elements in Human and Animal Nutrition, Fourth ed. Academic Press, New York. p. 170-196.

Ungerstedt, U. 1971. On the anatomy, pharmacology and function of the nigro-striatal dopamine system. Acta Physiol. Scand. Suppl. 367.

U.S. Bureau of Mines. 1983. Monthly Industry Survey, December, 1982. U.S. Dept of Interior, Washington, DC.

U.S. DHEW (U.S. Department of Health, Education and Welfare). 1958. Air pollution measurements of the national air sampling networks. Analysis of suspended particulate samples collected 1953-1957. PHS, Bur. State Serv. Div. Sanit. Eng. Serv., Robert A. Taft Sanitary Engineering Center, Cincinnati, OH. PHS Publ. 637. Washington, DC. 259 p.

U.S. DHEW (U.S. Department of Health, Education and Welfare). 1962. Motor vehicles, air pollution, and health. A report of the Surgeon General to the U.S. Congress. PHS, Div. Air Pollut., Washington, DC. p. 274-281.

U.S. DHEW (U.S. Department of Health, Education and Welfare). 1967. Parkersburg, West Virginia - Marietta, OH air pollution abatement activity. PHS, Natl. Center for Air Pollut. Control. March.

U.S. DHEW (U.S. Department of Health, Education and Welfare). 1970. Community water supply study: Analysis of national survey findings. PHS, U.S. DHEW, Environ. Health Serv., Bur. Water Hyg., Washington, DC. 123 p. PB 214 982; PB 215 198.

U.S. DHEW (U.S. Department of Health, Education and Welfare). 1978. Plan and operation of the HANES I Augmentation Survey of Adults 25-74 years. DHEW Pub. No. (PHS) 78-1314. Public Health Service. Natl. Center for Health Stat., Hyattsville, MD. June.

Usdin, T.B., I. Creese and S.H. Snyder. 1980. Regulation by cations of [ $^3\text{H}$ ]spiroperidol binding associated with dopamine receptors of rat brain. J. Neurochem. 34: 669-676.

U.S. EPA. 1971. National inventory of sources and emissions: Manganese - 1966. NTIS PB 220 020. Office of Air Quality Planning and Standards, U.S. EPA, Research Triangle Park, NC. August.

U.S. EPA. 1973. Handbook for Monitoring Industrial Wastewater. U.S. EPA, Technology Transfer, August.

U.S. EPA. 1974. Air pollution control engineering and cost study of the ferroalloy industry. NTIS PB 236 762. Control Programs Development Division, Office of Air Quality Planning and Standards, U.S. EPA, Research Triangle Park, NC. May.

U.S. EPA. 1975. Scientific and Technical Assessment Report on Manganese. EPA 600/6-75-002. ORD, Washington, DC. PB 242 291.

U.S. EPA. 1976. Quality Criteria for Water. GPO-1977-0-222-904. Washington, DC. 256 p.



U.S. EPA. 1977a. National trends in trace metals in ambient air 1965-1974. NTIS PB 264 906. Office of Air Quality Planning and Standards, Research Triangle Park, NC. February.

U.S. EPA. 1977b. A summary of industry comments received on manganese fuel additive (HMT). NTIS PB 123515. Office of Mobile Source Air Pollution Control, Technology Assessment and Evaluation Branch, U.S. EPA, Ann Arbor, MI

U.S. EPA. 1978a. Industrial Guide for Air Pollution Control. EPA 625/6-78-004. Technology Transfer, June.

U.S. EPA. 1978b. The multielemental analysis of drinking water using proton-induced X-ray emission (PIXE). EPA-600/1-78-058. Health Effects Research Laboratory. Cincinnati, OH. September.

U.S. EPA. 1979a. Air quality data for metals 1976 from the National Air Surveillance Networks. EPA-600/4-79-054. Environ. Monitoring and Support Lab. Research Triangle Park, NC. August.

U.S. EPA. 1979b. Ferroalloy process emissions measurement. NTIS PB 293 171. Industrial Environmental Research Laboratory, Office of Energy, Minerals and Industry. U.S. EPA, Research Triangle Park, NC. February.

U.S. EPA. 1979c. The Carcinogen Assessment Group's Preliminary Risk Assessment on Manganese. Washington, DC.

U.S. EPA. 1930. Guidelines and methodology used in the preparation of health effects assessment chapters of the consent decree water quality criteria. Fed. Reg. 45, 79347-79357.

U.S. EPA. 1981a. Receptor model technical series. Vol. 2. Chemical mass balance. NTIS PB 82 187345. Office of Air Quality Planning and Standards, Monitoring and Data Analysis Div. Research Triangle Park, NC.

U.S. EPA. 1981b. Level I environmental assessment of electric submerged-arc furnaces producing ferroalloys. NTIS PB 81-210106. Industrial Environmental Research Laboratory, Office of Environmental Engineering and Technology, U.S. EPA, Research Triangle Park, NC. March.

U.S. EPA. 1981c. Receptor models relating ambient suspended particulate matter to sources. NTIS PB 82-189549. Industrial Environmental Research Laboratory, U.S. EPA, Research Triangle Park, NC. March.

U.S. EPA. 1982. Review of the National Ambient Air Quality Standards for Particulate Matter: Assessment of Scientific and Technical Information. EPA-450/5-82-001. Office of Air Quality Planning and Standards, U.S. EPA, Research Triangle Park, NC. January.

U.S. EPA. 1983a. Ultrasonic extraction with heated  $\text{HNO}_3$ -HCl. Internal Standard Operating Procedure No. EMSL/RTP-SOP-EMD-003. Undergoing Agency review. Trace Element Analysis Section, Pollutant Analysis Branch, Environmental Monitoring Division, Environmental Monitoring and Support Laboratory, U.S. EPA, Research Triangle Park, NC.

U.S. EPA. 1983b. Inductively coupled argon plasma optical emission spectrometry. Internal Standard Operating Procedure No. EMSL/RTP-SOP-EMD-002. Undergoing Agency review. Trace Element Analysis Section, Pollutant Analysis Branch, Environmental Monitoring Division, Environmental Monitoring and Support Laboratory. U.S. EPA, Research Triangle Park, NC.

U.S. FDA (U.S. Food and Drug Administration). 1978. Compliance program evaluation. FY76 selected minerals in foods survey -- Adults, infants and toddlers (7320.63 and 7320.59). Industry Programs Branch, Bureau of Foods, U.S. FDA, Washington, DC. May 30.

U.S. PHS. 1962. Drinking Water Standards. U.S. DHEW, PHS Publ. No. 956, Washington, DC. In: National Academy of Sciences, 1977. Drinking Water and Health, NAS, Washington, DC.

Utter, M.F. 1976. The biochemistry of manganese. Med. Clin. North Am. 60: 713-727.

Van Beukering, J.A. 1966. Incidence of pneumonia among miners in an iron mine and a manganese mine in South Africa. Ned. Tijdschr. Geneesk. 110: 473-474.

Vanderborght, B.M. and R.E. Van Grieken. 1977. Enrichment of trace metals in water by adsorption on activated carbon. Anal. Chem. 49: 311-316.

Van Ormer, D.G. and W.C. Purdy. 1973. The determination of manganese in urine by atomic absorption spectrometry. Anal. Chim. Acta. 64: 93-105.

Versteck, J. and R. Cornelis. 1980. Normal levels of trace elements in human blood plasma or serum. *Anal. Chim. Acta.* 116: 217-254.

Versteck, J., A. Speecke, J. Hoste and F. Barbier. 1973. Determination of manganese, copper and zinc in serum and packed blood cells by neutron activation analysis. *Z. Klin. Chem. Klin. Biochem.* 11: 193-196.

Versteck, J., F. Barbier, A. Speeko and J. Hostle. 1974. Manganese, copper and zinc concentrations in serum and packed blood cells during acute hepatitis, chronic hepatitis and post-hepatic cirrhosis. *Clin. Chem.* 20: 1141-1145.

Vigliani, E. 1937. Esiste una polmonite de manganese? *Folia Med. (Naples).* 23: 451-458. (Ita.)

Volchok, H.L. and D.C. Bogen. 1973. Atmospheric trace metal studies. *In: Progress in Analytical Chemistry, Vol. 5, S. Ajuja et al., Ed. Chemical Analysis of the Environment and other Modern Techniques. Plenum Press, New York.* p. 17-41.

von Ottingen, W.F. 1935. Manganese: Its distribution, pharmacology and health hazards. *Physiol. Rev.* 15: 175-201.

Voss, H. 1939. Über das Vorkommen von beruflichen Manganvergiftungen in der Stahlindustrie (anlässlich eines Falles von Manganismus bei einem Ferromanganmüller). *Arch. Gewerbepath. Gewerbehyg.* 9: 453-463.

Voss, H. 1941. Spinal cord and peripheral nervous system in chronic manganese poisoning. Arch. Gewerbepath. Gewerbehyg. 10: 550-568.

Youk, V.B. and W.T. Piver. 1983. Metallic elements in fossil fuel combustion products. Environ. Health Perspect. p. 47. (In press)

Walton, K.G. and R.J. Baldessarini. 1976. Effects of  $Mn^{2+}$  and other divalent cations on adenylate cyclase activity in rat brain. J. Neurochem. 27: 557-564.

Wassermann, D. and M. Wassermann. 1977. The ultrastructure of the liver cell in subacute manganese administration. Environ. Res. 14: 379-390.

Wassermann, M. and G. Mihail. 1961. Recherches concernant la pathologie professionnelle des mineurs des mines de manganese. Arch. Gewerbepath. Gewerbehyg. 18: 632-657.

Wassermann, M. and G. Mihail. 1964. Indicateurs significatifs pour le depistage precoce du manganisme chez les mineurs des mines de manganese. Acta Med. Leg. Soc. 17: 61-89. (Fre.)

Wassermann, M., V. Voiculescu, A. Polingher, et al. 1954. Contribution to the understanding and prevention of manganism in manganese mines in the People's Republic of Rumania. Academia Republicii Populare Romine. Filiala Iasi. Studii si Ceretari Stiintifice. Seria 2, Stiinte Biologice, Medicale si Agricole. 5: 213-224.

Watanabe, H., S. Berman and D.S. Russell. 1972. Determination of trace metals in water using X-ray fluorescence spectrometry. *Talanta*. 19: 1363-1375.

Waters, M.D., D.E. Gardner, C. Aranyi and D.L. Coffin. 1975. Metal toxicity for rabbit alveolar macrophages in vitro. *Environ. Res.* 9: 32-47.

Weast, R.C. 1980. Handbook of Chemistry and Physics, 61st ed. The Chemical Rubber Co., Cleveland, OH. p. B24-25, B117, F24.

Wenlock, R.W., D.H. Buss and E.J. Dixon. 1979. Trace nutrients. 2. Manganese in British food. *J. Nutr.* 41: 253-261.

Westman, N.G. and S.L. Marklund. 1981. Copper- and zinc-containing superoxide dismutase in human tissues and human malignant tumors. *Cancer Res.* 41: 2962-2966.

WHO (World Health Organization). 1970. European Standards for Drinking Water, Second ed., Geneva, Switzerland. In: National Academy of Sciences, 1977. Drinking Water and Health, NAS, Washington, DC.

WHO (World Health Organization). 1976. Selected Methods of Measuring Air Pollutants. Offset Publ. No. 24. Geneva. p. 3-9.

WHO (World Health Organization). 1980. Recommended Health-based Limits in Occupational Exposure to Heavy Metals: Manganese. Report of a WHO Study Group. Tech. Rep. Ser. 647, WHO, Geneva, Switzerland. p. 80-101.

WHO (World Health Organization). 1981. Environmental Health Criteria 17. Manganese. WHO, Geneva.

Widdowson, E.M. 1969. Trace elements in human development. In: Mineral Metabolism in Paediatrics, D. Barltrop and W.L. Burland, Ed. Blackwell Sci. Publ., Oxford. p. 85-98.

Widdowson, E.M., H. Chan, G.E. Harrison and R.D.G. Milner. 1972. Accumulation of Cu, Zn, Mn, Cr and Co in the human liver before birth. Biol. Neonate. 20: 360-367.

Webe, A.H. 1930. The manganese content of the Mississippi River water at Fairport, IA. Science. 71: 248.

Wilgus, H.S. and A.R. Patton. 1939. Factors affecting manganese utilization in the chicken. J. Nutr. 18: 35-45.

Wilgus, H.S., L.C. Norris and G.F. Heuser. 1936. The role of certain inorganic elements in the cause and prevention of perosis. Science. 84: 252-253.

Witschi, H.P., P.J. Hakkinen and J.P. Kehrer. 1981. Modification of lung tumor development in A/J mice. Toxicology. 21: 37-45.

Witzleben, C.L. 1969. Manganese-induced cholestasis: Concurrent observations on bile flow rat and hepatic ultrastructure. Am. J. Pathol. 57(3): 617-626.

Witzleben, C.L. 1971. Bilirubin as a cholestatic agent: Physiologic and morphologic observations. *Am. J. Pathol.* 62: 181-194.

Witzleben, C.L. 1972. Physiologic and morphologic natural history of a model of intrahepatic cholestasis. *Am. J. Pathol.* 66: 577-588.

Witzleben, C.L., P. Pitlick, J. Bergmeyer and R. Benoit. 1968. Acute manganese overload. A new experimental model of intrahepatic cholestasis. *Am. J. Pathol.* 53: 409-423.

Wong, N.P., D.E. LaCroix and J.A. Alford. 1978. Mineral content of dairy products. II. Cheeses. *J. Am. Diet. Assoc.* 72: 608-611.

Wright, J.R., R. Levick and H.J. Atkinson. 1955. Trace element distribution in virgin profiles representing four great soil groups. *Proc. Soil Sci. Soc. Am.* 19: 3-44.

Wright, W.E., G.L. Ter Haar and E.B. Rifkin. 1973. The effect of manganese on the oxidation of  $\text{SO}_2$  in the air. Ethyl Corporation Research Laboratories, Detroit, MI. Nov.

Wundt, K., H. Duschner and K. Starke. 1979. Target preparation for X-ray emission analysis by anodic electrodeposition of cyano metalates from 2-propanol-water mixtures. *Anal. Chem.* 51: 1487-1492.

Wurts, T.C. 1959. Industrial sources of air pollution-metallurgical. In: *Proc. Natl. Conf. Air Pollut.*, Nov. 18-20, 1958. PHS Publ. 654. U.S. OHEW, p. 161-164.



Yamamoto, H. and T. Suzuki. 1969. Chemical structure of manganese compounds and their biological effects. In: Proceedings of the 42nd Annual Meeting of the Japan Association of Industrial Health, 28-31 March 1969. Jap. Assoc. Ind. Health, Fukuoka City.

Zaidi, S.H., R.K.S. Dogra, R. Shanker and S.V. Chandra. 1973. Experimental infective manganese pneumoconiosis in guinea pigs. Environ. Res. 6: 287-297.

Zhernakova, T.V. 1967. Correlation between iron, manganese and copper content in the blood serum of healthy individuals. Bull. Exp. Biol. Med. 63: 47-48.

Zielhuis, R.L., P. del Castillo, R.E.M. Herber and A.A.E. Wibowo. 1978. Levels of lead and other metals in human blood: Suggestive relationships, determining factors. Environ. Health Perspect. 25: 103-109.

## APPENDIX

## ESTIMATING HUMAN EQUIVALENT INTAKE LEVELS FROM ANIMAL STUDIES

## TERMINOLOGY AND APPROACH

The quantitative evaluation of potential health hazards for noncarcinogenic toxicants is based upon estimates of the threshold exposure level for the critical effect. Exposure levels for each study are evaluated as follows:

- NOEL No-Observed-Effect Level: That exposure level at which there are no statistically significant increases in frequency or severity of effects between the exposed population and its appropriate control.
- NOAEL No-Observed-Adverse-Effect Level: That exposure level at which there are no statistically significant increases in frequency or severity of adverse effects between the exposed population and its appropriate control. Effects are produced at this level, but they are not considered to be adverse.
- LOAEL Lowest-Observed-Adverse-Effect Level: The lowest exposure level in a study or group of studies which produces statistically significant increases in frequency or severity of adverse effects between the exposed population and its appropriate control.
- FEL Frank-Effect Level: That exposure level which produces unmistakable adverse effects, ranging from reversible histopathological damage to irreversible functional impairment or mortality, at a statistically significant increase in frequency or severity between an exposed population and its appropriate control.

The threshold estimate is bracketed by the highest NOEL and the LOAEL. The values for the NOELs and LOAEL depend on which health effect is considered. The estimate of the human threshold level is more uncertain if based on data for animals rather than for humans since there is presently limited information on species differences regarding toxic responses. Nevertheless, given limited dose-response data for humans it is necessary to extrapolate from the animal data.

Human equivalent intake rate (HEI) is defined here as the exposure level estimated from animal data which would cause the same health effect in humans if continued over the same fraction of lifespan as used in the animal

study. The conversion for manganese assumes that if the ratio (exposure level)/(body surface area) is the same in humans as in the animal study, then effects of the same severity will occur.

#### CRITICAL EFFECTS AND ESTIMATED EFFECT LEVELS

The lowest exposure level for humans associated with adverse effects is an estimated LOAEL of 3-11  $\mu\text{g}/\text{m}^3$  (based on emissions from a ferromanganese plant) for respiratory effects in children reported by Nogawa et al. (1973). Comparison among studies of respiratory effects in laboratory animals (summarized in Table 9-1) shows that Ulrich et al. (1979a,b,c) and Suzuki et al. (1978) utilized the longest exposure periods at exposure concentrations  $\sim 100 \mu\text{g}/\text{m}^3$ . Although there are shortcomings in each study (see Section 9.5.) the repeated observations suggest that this level may be close to the threshold. Therefore, these studies were selected for these risk assessment calculations. The HEI is estimated from the data from experimental animals by the following:

$$\text{HEI} = C_A \times D_E \times \text{Br} \times \left( \frac{70 \text{ kg}}{W_a} \right)^{2/3}$$

where  $C_A$  = concentration in air in the animal study in  $\mu\text{g}/\text{day}$

$D_E$  = fraction of day experimental animals were exposed

$\text{Br}$  = volume of air breathed per day in  $\text{m}^3$

$W_a$  = body weight of the experimental animal in kg

This conversion is based on the following assumptions:

1. Agents that are in the form of particulate matter are expected to be absorbed and retained proportional to the breathing rate.
2. The fraction retained is approximately the same for all species.
3. The conversion from animals to humans based on exposure level per body surface area more accurately reflects differences among species than does a mg/kg body weight conversion (Rall, 1969). The surface area ratio is well approximated by the body weight ratio to the 2/3 power (Calabrese, 1983).

The estimation of HEI is based on intake by inhalation of manganese in excess of dietary intake of this essential element because all studies used here were so designed. Also, a sufficient oral intake and strong homeostatic control can be assumed so that excess exposure is appropriate. Inhalation studies should be used since the critical effect is route specific.

The estimated HEI (in mg/day) is converted to a human equivalent exposure level (HEEL) by dividing by the average daily human respiratory rate of  $20 \text{ m}^3/\text{day}$ . All calculations are summarized in Table A-1.

In the studies considered here  $D_e$  equals 1. Therefore, for rhesus monkeys in the study by Suzuki et al. (1978)

$$\begin{aligned} \text{HEI} &= C_A \times \text{Br} \times \left( \frac{70 \text{ kg}}{W_a} \right)^{2/3} \\ &= 700 \text{ } \mu\text{g}/\text{m}^3 \times 1.4 \text{ m}^3/\text{day} \times \left( \frac{70 \text{ kg}}{3.5 \text{ kg}} \right)^{2/3} \\ &= 7293 \text{ } \mu\text{g}/\text{day}. \end{aligned}$$

The HEEL obtained by dividing by the daily respiratory volume ( $20 \text{ m}^3/\text{day}$ ) is  $365 \text{ } \mu\text{g}/\text{m}^3$ . Note that this HEEL is based on a LOAEL and thus may be above the human threshold level.

Similar calculations using data on rats from Ulrich et al. (1979a,b,c) ( $W_a = 0.35 \text{ kg}$ ,  $\text{Br} = 0.26 \text{ m}^3/\text{day}$ , and  $C_A = 113 \text{ } \mu\text{g}/\text{m}^3$ ) result in a HEEL of  $51 \text{ } \mu\text{g}/\text{m}^3$ . For the Ulrich et al. (1979a,b,c) data on squirrel monkeys ( $W_a = 0.72 \text{ kg}$ ,  $\text{Br} = 0.72 \text{ m}^3/\text{day}$ , and  $C_A = 113 \text{ } \mu\text{g}/\text{m}^3$ ) the HEEL is  $87 \text{ } \mu\text{g}/\text{m}^3$ .

These results, shown in Table A-1, can be compared with the data from Nogawa et al. (1973) on children who had an estimated LOAEL of  $3\text{--}11 \text{ } \mu\text{g}/\text{m}^3$ . The data obtained from human data is, of course, crucial for public health decision making. This level might be expected to be lower than the animal level for the following reasons: 1) the studies on animals

TABLE A-1

Exposure Effect Information for Health Hazard Evaluation: Human Equivalent Exposure Levels Estimated from Animal Data

Species	Response Level	Exposure ( $\mu\text{g}/\text{m}^3$ )	<u>Animal Intake Level</u>		<u>Estimated Human Intake Level</u>		Estimated Human Equivalent Exposure Level (HEEL) ( $\mu\text{g}/\text{m}^3$ )	Adjusted HEEL <sup>a</sup>	Reference
			$\mu\text{g}/\text{day}$	$\mu\text{g}/\text{kg}/\text{day}$	$\mu\text{g}/\text{day}$	$\mu\text{g}/\text{kg}/\text{day}$			
A-4 Rat	reported NOEL <sup>b</sup>	113	29	84	1022	15	51	5	Ulrich et al., 1979a,b,c
Squirrel monkey	reported NOEL <sup>b</sup>	113	81	113	1740	25	87	8.7	Ulrich et al., 1979a,b,c
Rhesus monkey	LOAEL	700	980	280	7293	104	365	36.5	Suzuki et al., 1978

<sup>a</sup>The estimated HEEL is adjusted by dividing by an uncertainty factor of 10 to compensate for the heterogeneity in human populations and thereby protect the sensitive individual.

<sup>b</sup>These NOELs are used to compare human and animal data despite some limitations which exist in the study. See text for further discussion.

have flaws resulting in uncertainty as to whether adverse effects are missed; 2) certain endpoints studied in humans cannot be ascertained as well in animal studies and are likely to be overlooked; and 3) children are a sensitive group. A better comparison between animal and human data is obtained by dividing the HEEL by 10 to compensate for the heterogeneity in the human population and to better protect the sensitive individual (Dourson and Stara, 1983). The range of adjusted HEELs from the animal NOELs and the LOAEL is 5-37  $\mu\text{g}/\text{m}^3$ , and supports the human LOAEL of 3-11  $\mu\text{g}/\text{m}^3$  observed in a sensitive population.

END

DATE

FILMED

3-12-87

NTIS

# TABLE OF CONTENTS

Executive Summary .....	1
Introduction .....	3
HEI Involvement .....	5
Framework and Process .....	6
Background Information on Manganese .....	8
Ambient Levels of Manganese .....	8
Dietary Intake and Body Burden .....	9
Health Effects of Manganese .....	11
Modifying Factors .....	11
Fate of Inhaled Manganese .....	15
Neurotoxic Effects .....	17
Pulmonary Toxicity .....	20
Exposure to Manganese .....	27
Anticipated Exposure Levels from	
Automotive Sources .....	27
Characteristics of Manganese Emission .....	30
Determination of the Inhaled	
Dose of Manganese .....	31
Discussion and Conclusions .....	33
Estimation of the Worst-Case	
Inhaled Dose .....	33
Potential Neurotoxic Effects .....	35



## EXECUTIVE SUMMARY

The use of ceramic particulate traps, in conjunction with manganese fuel additives, has been viewed as a way to reduce emissions of particulate matter from diesel-fueled vehicles. This report focuses on the potential health effects from increased public exposure to manganese emissions from such a use.

Manganese is the 12th most abundant element on earth. Ambient levels of manganese rarely exceed  $0.1 \mu\text{g}/\text{m}^3$ . Manganese is classified as an essential element for physiologic functions. Average daily intake of manganese, which is mostly from food, is about 3 to 4 mg/day (range 2 to 9 mg/day). Homeostatic mechanisms (primarily by regulation of intestinal absorption and biliary excretion) maintain the body burden of manganese between 12 and 20 mg.

The health effects of major concern from manganese exposure are neurotoxic and respiratory. Numerous studies, mostly in occupational groups exposed to very high levels of manganese (in the  $\text{mg}/\text{m}^3$  range), have demonstrated that prolonged exposure to relatively high levels of manganese leads to manganism (a Parkinson-disease like syndrome) and other symptoms of central nervous system toxicity, as well as to pneumonitis, bronchitis, and increased susceptibility to pulmonary infections. Pulmonary effects of manganese have been observed to occur at lower levels of exposure than neurotoxic effects.

The health effects of exposure to low levels of manganese have not been well characterized. An epidemiologic

study (Saric et al. 1977) has reported neurotoxic effects of manganese at exposure levels that were between 300 and 5,000  $\mu\text{g Mn/m}^3$ . Another epidemiologic study (Nogawa et al. 1973), which focused on respiratory symptoms in children attending a school near a ferromanganese plant, has reported health effects at levels as low as 3 to 11  $\mu\text{g Mn/m}^3$ . However, serious problems with the study (such as poor characterization of exposure, failure to blind observers and to randomize students, and a lack of control for the confounding factors) suggest that the results of the study are not reliable. All other studies have reported adverse effects at levels that are one to two orders of magnitude greater.

A modeling study by Volkswagen (1984) for emissions of manganese suggests that the worst-case ambient concentration of manganese in an urban street canyon is likely to be less than 0.5  $\mu\text{g/m}^3$  (the assumptions include 6.6 percent penetration of the manganese technology into the vehicle fleet, emissions of 100 percent of the manganese added to the fuel in the emissions, high manganese consumption, and worst-case traffic conditions). HEI has independently verified the Volkswagen calculations. If an individual, such as a traffic policeman, worked outdoors all day in a street with the worst-case levels of manganese, his daily intake of manganese by inhalation from ambient sources would be 3.6  $\mu\text{g}$ . This is a very small amount compared to the daily dietary intake of manganese (3 to 4 mg/day), and is not expected to tax the body's homeostatic mechanisms that regulate manganese.

On the basis of the information in the literature, and the exposure and dose of manganese estimated in this report and summarized above, it appears very unlikely that exposure to airborne manganese from mobile sources (worst-case level 0.5  $\mu\text{g/m}^3$ ) would produce adverse neurologic effects.

Regarding respiratory effects, the difference between estimated exposure (worst-case level  $0.5 \mu\text{g Mn/m}^3$ ) and the lowest level at which health effects have been reported in the Nogawa et al. (1973) study ( $3$  to  $11 \mu\text{g Mn/m}^3$ ) is not very large. However, in view of the concerns regarding the Nogawa study, it appears unlikely that the increased ambient levels of manganese from trap-equipped diesel vehicles would produce adverse respiratory effects.

Given the paucity of information at the anticipated low levels of manganese from its use in diesel-powered vehicles, if manganese were to be used as a fuel additive, it is very important that the emissions of manganese be characterized in terms of their physical form and chemical species, the assumptions made in estimating the emissions, exposure, and dose of manganese be substantiated, and the results of the Nogawa study be reviewed and evaluated in greater depth than attempted in this report.

### INTRODUCTION

Combustion of diesel fuel results in the formation of fine carbonaceous particles, a number of organic compounds that can adsorb to the particles, and a number of gases. Because of concerns with the health effects of diesel emissions, the U.S. Environmental Protection Agency (EPA) has promulgated regulations that are aimed at reducing the emission of particulate matter from diesel-powered vehicles. The EPA has also mandated that the emissions of nitrogen oxides from diesel-powered automobiles be reduced. However, because of the nature of the combustion process, technologies for reducing particulate emissions tend to increase emissions of oxides of nitrogen, and vice versa. The general approach adopted in the automobile industry to meet these regulations